

Protocol

The Cost Effectiveness of a Tailored, Web-Based Care Program to Enhance Postoperative Recovery in Gynecologic Patients in Comparison With Usual Care: Protocol of a Stepped Wedge Cluster Randomized Controlled Trial

Esther VA Bouwsma^{1,2,3}, MD; Johannes R Anema^{2,3}, MD, PhD; Antonie Vonk Noordegraaf^{1,2,3}, MD; Dirk L Knol⁴, PhD; Judith E Bosmans^{3,5}, PhD; Steven E Schraffordt Koops⁶, MD, PhD; Paul JM van Kesteren⁷, MD, PhD; W Marchien van Baal⁸, MD, PhD; Jos P Lips⁹, MD, PhD; Mark H Emanuel¹⁰, MD, PhD; Petrus C Scholten¹¹, MD, PhD; Alexander Mozes¹², MD; Albert H Adriaanse¹³, MD, PhD; Hans AM Brölmann^{1,3}, MD, PhD; Judith AF Huirne^{1,3}, MD, PhD

¹Department of Obstetrics and Gynecology, VU University Medical Center, Amsterdam, Netherlands

²Department of Public and Occupational Health, VU University Medical Center, Amsterdam, Netherlands

³EMGO Institute for Health and Care Research, Amsterdam, Netherlands

⁴Department of Epidemiology and Statistics, VU University Medical Center, Amsterdam, Netherlands

⁵Department of Health Sciences, Faculty of Earth and Life Sciences, VU University, Amsterdam, Netherlands

⁶Department of Obstetrics and Gynecology, Meander Medical Center, Amersfoort, Netherlands

⁷Department of Obstetrics and Gynecology, Onze Lieve Vrouwe Gasthuis, Amsterdam, Netherlands

⁸Department of Obstetrics and Gynecology, Flevo Hospital, Almere, Netherlands

⁹Department of Obstetrics and Gynecology, Kennemer Gasthuis, Haarlem, Netherlands

¹⁰Department of Obstetrics and Gynaecology, Spaarne Hospital, Hoofddorp, Netherlands

¹¹Department of Obstetrics and Gynaecology, Diakonessenhuis, Utrecht, Netherlands

¹²Department of Obstetrics and Gynecology, Amstelland Hospital, Amstelveen, Netherlands

¹³Department of Obstetrics and Gynecology, Medical Center Alkmaar, Alkmaar, Netherlands

Corresponding Author:

Judith AF Huirne, MD, PhD

Department of Obstetrics and Gynecology

VU University Medical Center

De Boelelaan 1117

Amsterdam,

Netherlands

Phone: 31 20444 4827

Fax: 31 204444811

Email: j.huirne@vumc.nl

Abstract

Background: The length of recovery after benign gynecological surgery and return to work frequently exceeds the period that is recommended or expected by specialists. A prolonged recovery is associated with a poorer quality of life. In addition, costs due to prolonged sick leave following gynecological surgery cause a significant financial burden on society.

Objective: The objective of our study was to present the protocol of a stepped wedge cluster randomized controlled trial to evaluate the cost effectiveness of a new care program for patients undergoing hysterectomy and/or adnexal surgery for benign disease, compared to the usual care.

Methods: The care program under study, designed to improve convalescence and to prevent delayed return to work, targets two levels. At the hospital level, guidelines will be distributed among clinical staff in order to stimulate evidence-based patient education. At the patient level, additional perioperative guidance is provided by means of an eHealth intervention, equipping patients with tailored convalescence advice, and an occupational intervention is available for those patients at risk of prolonged sick leave. Due to the stepped wedge design of the trial, the care program will be sequentially rolled out among the 9 participating

hospitals, from which the patients are recruited. Eligible for this study are employed women, 18-65 years of age, who are scheduled for hysterectomy and/or laparoscopic adnexal surgery. The primary outcome is full sustainable return to work. The secondary outcomes include general recovery, quality of life, self-efficacy, coping, and pain. The data will be collected by means of self-reported electronic questionnaires before surgery and at 2, 6, 12, 26, and 52 weeks after surgery. Sick leave and cost data are measured by monthly sick leave calendars, and cost diaries during the 12 month follow-up period. The economic evaluation will be performed from the societal perspective. All statistical analyses will be conducted according to the intention-to-treat principle.

Results: The enrollment of the patients started October 2011. The follow-up period will be completed in August 2014. Data cleaning or analysis has not begun as of this article's submission.

Conclusions: We hypothesize the care program to be effective by means of improving convalescence and reducing costs associated with productivity losses following gynecological surgery. The results of this study will enable health care policy makers to decide about future implementation of this care program on a broad scale.

Trial Registration: Netherlands Trial Register: NTR2933; <http://www.trialregister.nl/trialreg/admin/rctview.asp?TC=2933> (Archived by WebCite at <http://www.webcitation.org/6Q7exPG84>).

(*JMIR Res Protoc* 2014;3(2):e30) doi: [10.2196/resprot.3236](https://doi.org/10.2196/resprot.3236)

KEYWORDS

gynecology; Internet; telemedicine; convalescence; return to work; economic evaluation

Introduction

Early Discharge From the Hospital

In the last two decades, the hospital stay following surgical procedures has been shortened drastically, due to recovery-enhancing strategies such as the use of minimally invasive techniques and the implementation of fast-track programs [1-4]. The advantages of early postoperative discharge include increased patient satisfaction, low hospital-acquired infection rates, and reduced hospitalization costs [5]. However, a major disadvantage of minimizing the length of a hospitalization is that patient contact becomes very brief, which is often at the expense of time spent on patient education. Ironically, the lack of detailed convalescence instructions at the time of discharge increases the risk of an unnecessary prolonged recovery [6-11]. Therefore, as long as the organization of perioperative care has not fully anticipated the transition of postoperative recovery to the home setting, early discharge does not necessarily translate into accelerated recovery and earlier resumption of (work) activities [12-14].

In gynecology, the postoperative convalescence after discharge from the hospital has not received much attention in research and practice. Yet, there is considerable evidence that the length of recovery time after a gynecological surgery systematically exceeds the period considered as appropriate by specialists [5,10,12-17]. In a prospective study performed by our own study group among 148 patients receiving gynecological surgery for a benign disease, median time to return to work (RTW) exceeded the recommended sick leave of 6 weeks by approximately 3 weeks. The median time to RTW following an intermediate surgery (eg, laparoscopic or vaginal hysterectomy) was 60 days (interquartile range, IQR 56-135) and following a major surgery (eg, abdominal hysterectomy) 69 days (IQR 56-135) [10].

Prolonged Recovery at Home

An unnecessary prolonged recovery is associated with poorer quality of life [18,19]. In addition, work related problems have

also been associated with an increase in health care consumption [20]. Furthermore, taken into account that about 14,000 hysterectomies are performed annually in the Netherlands alone [21], the financial burden on society due to delayed convalescence after a gynecological surgery is substantial.

In order to reduce unnecessary delayed recovery, and concurrently decrease costs associated with prolonged sick leave and increased health care utilization following gynecological surgery, our research group started working on an innovative strategy to optimize perioperative care in 2008. Since the beginning of the project several goals were achieved, starting with the development of detailed convalescence recommendations following 4 types of benign gynecological surgery, using a modified Delphi method [22]. Simultaneously, a multidisciplinary care program was developed [23,24] consisting of an interactive eHealth intervention and—for those patients at risk of prolonged sick leave—an occupational intervention. The care program provides guidance to patients from the moment the surgery is planned, until the full resumption of all activities—including return to work—and encourages patients to take an active role in their own recovery. The care program was subject to an effect evaluation as well as a process evaluation in 2010 [25]. While the effectiveness study among 215 patients showed a positive effect on the outcomes: (1) RTW, (2) quality of life, and (3) perceived pain [26], the process evaluation showed some room for improvement [27].

Besides evaluating the effectiveness of a study, it is of equal importance to conduct an economic evaluation, especially considering the high economic burden of extended time to convalescence after a gynecologic surgery. The economic evaluations are necessary to gain insight into the costs of an intervention in relation to its effects. Health care policy makers can use these results to decide how resources should optimally be allocated to maximize health or welfare [28].

Therefore, the primary objective of the current study is to conduct an economic evaluation of the care program compared to the usual care. This economic evaluation will be conducted

alongside a randomized trial, as the intervention concerns a further developed version of the care program, which has not yet been subject to an effect evaluation. In addition, this construction enables the systematic collection of relevant effect and cost data under “real life” conditions. As the intervention care program targets two levels (the hospital level and the patient level), a cluster design was chosen in order to prevent contamination between the study arms. The primary outcome duration until full sustainable RTW will be assessed on the level of the individual participant. On the level of the participating hospitals, we will investigate to what extent the guidelines on convalescence recommendations are adopted, and how future implementation of the guidelines and care program can be facilitated.

Methods

The Standard Protocol Items

The Standard Protocol Items, Recommendations for Interventional Trials statement [29], and CONSolidated Standards Of Reporting Trials (CONSORT) statement [30,31], were used in order to describe the design of this study. In addition, we used the extension to cluster randomized trials [32] and the CONSORT eHealth checklist [33].

Ethical Issues

The Institutional Review Boards of all participating hospitals approved this study protocol. Informed consent was obtained from all of the patients.


Trial Design

This trial is designed as a cluster, randomized controlled, stepped wedge trial, which involves a sequential rollout of the intervention in the participating clusters over several time

periods. In our study, clusters are the departments of obstetrics and gynecology in nine different hospitals in the Netherlands. Each time period (TP) takes 2 months. At the start of the trial (TP₁), all of the patients scheduled for a surgery in all of the participating hospitals receive usual care (control phase). After two months (TP₂), the intervention is implemented in the first cluster, and from now on the patients scheduled for a surgery in this hospital will receive the intervention program, while in all of the other hospitals the patients still receive usual care. The patients in cluster 2 who underwent surgery during TP₁ remain in the control group until they finish the 12 month follow-up. During TP₃, the intervention program continues in cluster 1, and the intervention is implemented in cluster 2 as well, resulting in the deliverance of the intervention program to the patients in clusters 1 and 2 that will undergo surgery from this point onward, while patients in clusters 3 to 9 serve as the control group. At the beginning of TP₄, cluster 3 starts with the intervention, etc. This is repeated until the intervention is implemented in all clusters (TP₁₀). Figure 1 illustrates the study design.

A cluster design was chosen to minimize the risk of contamination, as our intervention targets both health care providers and patients. A stepped wedge approach was employed because of the unique feature of an unidirectional crossover, preventing the intervention to be withdrawn from the hospital during the trial [34-36]. Because there is substantial evidence from our previous trial that the care program under study will be effective, this is particularly convenient, as hospitals will be able to keep using the intervention after the trial. Moreover, it enables us to study the implementation process carefully, giving valuable insight into barriers and facilitators for future broader implementation.

Figure 1. Trial design.



Selection of Clusters

The clusters in this trial consist of nine hospitals in the surroundings of Amsterdam, the capital of the Netherlands. The hospitals were eligible if they performed at least 100 hysterectomies or laparoscopic adnexal surgeries yearly, and were located within 50 km of the Vrije Universiteit Medical Center (VUmc). The research team enrolled the clusters before the start of the trial. In an attempt to select a heterogeneous sample of hospitals, we included 1 university hospital, 7 teaching hospitals, and 1 nonteaching hospital.

Study Population

The eligible participants for this study are women 18-65 years of age, employed for at least 8 hours per week (salary employed,

self employed, or voluntary work), and scheduled for a surgery for a benign gynecological disease in one of the nine participating hospitals. The types of surgeries that are included are: (1) total abdominal hysterectomy (TAH), (2) vaginal hysterectomy (VH), (3) total laparoscopic hysterectomy or laparoscopic assisted vaginal hysterectomy (TLH), or (4) laparoscopic adnexal surgery (LAS). The factors that are possibly complicating the postoperative course (eg, severe comorbidity, malignancy, pregnancy), the factors that are interfering with the eHealth intervention (computer or Internet illiteracy), or with the occupational intervention (conflict with employer, prolonged sick leave, or disability) serve as the exclusion criteria. [Table 1](#) lists an overview of all eligibility criteria.

Table 1. Eligibility criteria.

Inclusion criteria	Exclusion criteria
Women scheduled for:	(Suspicion of) malignancy
Laparoscopic adnexal surgery	(Ectopic) pregnancy
Total laparoscopic hysterectomy	Deep infiltrating endometriosis
Vaginal hysterectomy	Concomitant health problems affecting daily activities
Total abdominal hysterectomy	Psychiatric disorders affecting daily activities
18-65 years of age	Legal conflict with employer
Employed \geq 8 hours/week	Being sick listed $>$ 4 weeks, or when reason of sick leave is related to gynecological surgery $>$ 2 months
	Inability to understand or complete Dutch questionnaires
	Computer or Internet illiteracy

Recruitment of Patients

The recruitment of patients will take place in all participating hospitals. When the patients are scheduled for a hysterectomy or laparoscopic adnexal surgery, they will receive a letter about the study on behalf of their gynecologist. The letter includes detailed information about the trial. In addition, it is explained that someone from the research team will make contact by telephone after one week to evaluate the patients' willingness to participate and answer questions if necessary. If the patient does not wish to be contacted, she can return an included reply card, or send an email to a specified email address.

When contact is made and the patient is willing to participate, eligibility is assessed. The eligible patients are then requested to return a signed informed consent, which is also attached to the information letter. The participants will not receive any financial or nonfinancial incentives.

Randomization

The randomization takes place at the level of the clusters and determines the order in which the intervention program is implemented in the participating hospitals. The randomization will be performed by a statistician using a computer generated list of random numbers.

The patients are informed about the allocation of treatment by the research team after the patient's informed consent and the completion of the first questionnaire before surgery. As the treatment allocation depends on the scheduled date of the

surgery, and the implementation phase of the hospital in which they are being operated, it is predetermined for each participant, potentially causing selection bias. To minimize the risk of selection bias, the participants will not be informed about the study design, and will be counseled as if they have equal chances between receiving the usual care or the intervention program. For this reason, counseling will be done by the research team, rather than by their own physician, who might be, for example, more willing to include patients during the intervention phase than during the control phase. Moreover, physicians will be blinded to the randomization schedule, and will only be informed about the start of the intervention phase approximately one month before the actual implementation. Once the intervention phase has started, the importance of not communicating this information with the potential patients will be emphasized.

Interventions

Usual Care

Before the implementation of the intervention program, the participants receive the usual perioperative care as provided in the hospital in which they are scheduled for surgery. Although considerable variation exists in the Netherlands, in most cases patients get verbal (general) instructions at discharge by a nurse and/or physician, often followed—but not necessarily—by a letter or brochure. In general, an outpatient postoperative consultation is scheduled 4 to 6 weeks following the surgery. Between discharge and the postoperative consultation, medical

care is only initiated by the patient, who can consult her general physician (GP) or gynecologist, if necessary. Employed workers who have not resumed work within 6 weeks after the surgical procedure will be invited for a consultation with their occupational physician (OP), as required by law in the Netherlands.


Intervention

The systematic development of the care program using the principles of Intervention Mapping is described in more detail

elsewhere [23]. Both theory and practice were combined, and all stakeholders were involved in the process. The engagement of the patients was prompted through focus groups [24]. The Attitude, Social influence, and Self-efficacy model was used as a theoretical framework for determinants of behavior regarding return to work [37,38].

The care program targets two levels, which are described below. Figure 2 shows an overview of the intervention care program.

Figure 2. Overview of the care program. GP=general physician; OP= occupational physician; RTW=return to work.



Cluster Level

At the cluster level, the intervention care program aims to structure and stimulate evidence-based perioperative care. Approximately two months before a cluster shifts from the control to the intervention phase, the principle researcher will approach the head of the department to arrange logistics. A minimum of two meetings is planned one or two weeks before the actual implementation with physicians and nurses to provide and explain the new convalescence recommendations that should be communicated to the patients. In addition, all health professionals involved in the clinical care receive a pocket card on which these recommendations are summarized for quick reference. The residents involved in the discharge communication are instructed to explain the convalescence recommendations to their patients before they are discharged. Visual reminders in the patient records will help the residents do so. With the secretary of the department, a strategy is developed to prompt the standard postoperative consultation at 4 weeks following a hysterectomy, and 2 weeks following adnexal surgery. During the trial, newsletters will be spread regularly to reinforce the different aspects of the intervention care program.

Patient Level

Individual Tailored Guidance

At the patient level, the care program aims to provide individual tailored guidance to patients from the moment the surgery is planned until the full resumption of all activities. It consists of two steps: (1) access to an interactive eHealth intervention for all patients, and (2) an additional occupational intervention for those patients at risk for prolonged sick leave.

eHealth Intervention

The patient webportal ([Figure 3](#), [39]) aims at empowering its users and improving communication between patients and their employers, as well as improving the communication between the involved health care professionals during the perioperative period. Access to the webportal will be given to the patients approximately 2 to 4 weeks prior to surgery by the research team, by providing a username and temporary password. The instructions are given by email, and it is explained that if patients require assistance, they can contact the research team by phone

or email. If patients fail to log in, an automatic reminder is sent to them one week before their surgery to remind them about the webportal and its functionalities. User authentication will make it possible to analyze website activity for each individual participant (visit duration, number of sessions, number and details of pages visited).

The most important tool of the webportal is the possibility to generate a tailored convalescence plan. In the instruction email, patients are encouraged to generate such a plan at least once, preferably before surgery. Having access to detailed convalescence advice will enable the patients to develop realistic expectations about their own recovery, and plan the resumption of their activities and work reintegration accordingly. Moreover, a tailored convalescence plan will help the patients gain insight into potential recovery problems and find solutions at an early stage, preferably before surgery. Because the convalescence plan is composed before surgery, gynecologists are asked to approve the plan electronically on the first postoperative day. In the case of an uncomplicated procedure, the plan is turned into a definite convalescence plan, and the patients are instructed to follow the recommendations in it. In the case of a converted procedure, the plan is adjusted to the type of surgery that was actually performed. In the event of severe complications, the gynecologist can choose not to approve the convalescence plan, and the patients then receive a message that the convalescence plan is not valid anymore, and that they should follow up with the specific instructions given to them at discharge. With the consent of the patient, the approved convalescence plan is also disclosed to the GP and/or OP of the patient. This last feature was added since the prior evaluation of the webportal, and was developed to facilitate the involvement of other health care professionals during the perioperative period in order to stimulate a multidisciplinary approach. In addition, the webportal was equipped with a tool that enables the patients to generate a recovery report, a graphic presentation of their own recovery, allowing them to track their progress.

During the trial, the content of the website will be frozen, except from the dynamic component (forum). [Table 2](#) summarizes the most important tools of the eHealth intervention. Screenshots of the webportal are included as a Multimedia Appendix (see [Multimedia Appendix 1](#)).

Figure 3. Screenshot of ikherstel.

Table 2. Content of the eHealth intervention.

Tool	Description
Personalized convalescence plan ^a	The tool allows patients to generate detailed tailored instructions on the resumption of activities after the surgery, allowing preoperative planning of (work) activities. The convalescence plan is approved electronically on the first postoperative day by the surgeon who performed the surgery, resulting in a definitive convalescence plan. With the consent of the patient, the approved convalescence plan is shared with GP and/or OP. ^a
Recovery monitor + recovery report ^a	The tool makes an inventory of the resumption of activities at 2, 4, 7, 14, 28, 56, and 84 days after surgery. Results are graphically displayed in a recovery report, allowing the patient to track their progress. ^a In case the patients fall behind, an alerting system advises them to contact a specific health care professional, depending on the underlying problem.
Invitation of employer	The tool allows patients to invite an employer to an anonymous section of the webportal to stimulate a dialogue. The development of a reintegration plan preoperatively will help them gain insight into potential RTW problems.
Video	There is a 9-minute film illustrating the common pitfalls during the postoperative period.
Knowledge	There are several tools to find additional information, such as an extended list with answers to frequently asked questions, a glossary, and links to other useful websites.
Forum	The tool allows the patients to interact (privately or publicly) with other patients.

^aTools that were modified since the last evaluation of the webportal

Occupational Intervention

The occupational intervention is developed to provide additional guidance to those patients at risk for prolonged sick leave. The occupational intervention will be delivered by a group of six

independent OPs, who will be trained as RTW coordinators before the start of the trial. There are two types of consultations: (1) a preoperative, and (2) a postoperative consultation. All consultations will be delivered by telephone, unless the OP and the patient decide together otherwise.

The patients who have an inadequate expectation about their own recovery (longer than 3 weeks for LAS, longer than 6 weeks for VH/TLH, or longer than 8 weeks for TAH), or have a low intention to resume work activities while still recovering, are offered a preoperative consultation, as expectations about RTW and intention to resume work have been identified as two predictors for RTW in recent studies [10,40,41]. During the preoperative consultation, the OP explains the importance of a prosperous recovery in terms of improving quality of life and preventing long term sickness. In addition, the OP tries to identify and—if necessary—alter attitudes and (irrational) beliefs about recovery.

The patients who exceed 5 weeks of sick leave receive a postoperative consultation, during which, the OP assesses the underlying mechanism for the delayed recovery. The OP gives advice to improve the reintegration process. Moreover, as a RTW coordinator, the OP has an excellent position to communicate with the patient's gynecologist, GP, OP, and employer, if necessary, and of course, with the consent of the patient, stimulating an integrated care approach. In addition, the OP has the possibility to initiate a participatory workplace intervention, aimed at finding consensus between the patient and her employer concerning solutions for identified obstacles for RTW with the help of an occupational therapist (OT) [42,43].

The occupational intervention described above differs from the intervention as delivered during the first trial, due to the insight gained during the process evaluation. Originally, contact with the clinical OP took place in the 10th or 11th week, however, this turned out to be too late in order to be able to alter attitudes and beliefs, and influence the development of a solid RTW plan. Therefore, in the current trial, contact will be made much earlier, at 5 weeks, and on indication already before surgery. In addition, the patients will receive the details of the postoperative appointments before surgery in order to prepare them that the occupational intervention is part of the care program they receive, as in the prior trial, almost half of the patients declined additional occupational care. In the case of full RTW, the postoperative appointment will be cancelled.

Outcomes

Effect Measures

The effects of the intervention will be assessed on the level of the patient. The primary outcome of the study is the sick leave duration until full sustainable RTW, defined as the duration of the sick leave in calendar days from the day of surgery until full RTW, in their own work or other work with equal earnings, for at least 4 weeks without (partial or full) recurrence [44]. The recurrence of sick leave due to the gynecologic surgery within the four week period after initial full RTW will be added to the preceding period of the sick leave. The RTW will be assessed by a monthly electronic sick leave calendar.

Secondary outcomes that will be assessed are:

1. Recovery, measured by the Recovery Index-10 (RI-10) a validated recovery-specific questionnaire [45];

2. Self-reported quality of life, assessed by the Dutch versions of the EuroQol-5D (EQ-5D) [46] and the Short-Form Health Survey (SF-36) [47,48];
3. Duration of sick leave until first RTW, and total duration of sick leave due to the gynecological surgery for the entire follow-up period, both measured by the monthly sick leave calendars;
4. Self-efficacy, assessed by the Dutch adaptation of the General Self-Efficacy Scale (GSES) [49];
5. Coping, assessed by the Pearlin Mastery Scale (PMS) [50];
6. Pain intensity, measured by the Von Korff questionnaire (VAS) [51]; and
7. (Post) operative complications both assessed through self-report and by the review of surgical reports. Complications include: (1) enlargement of the wound (≥ 8 cm), (2) unintended injury to other structures (eg, bowel, bladder, ureter), (3) unexpected blood loss requiring transfusion, (4) prolonged hospital stay, (5) readmission within 72 hours (overnight), (6) repeat surgery within 2 weeks, and (7) postoperative infection requiring antibiotics.

Prognostic Factors

Before surgery, data about potential prognostic factors will be collected. In case of coincidental and meaningful differences, analyses will be adjusted for the following characteristics: (1) sociodemographic data such as age, education level, and ethnicity; (2) personal factors such as expectation, motivation, and intention toward RTW, duration of sick leave in the past 3 months; and (3) work related factors such as physical workload and potential work related psychosocial factors, assessed by the Dutch Musculoskeletal Questionnaire (DMQ) [52] and the Job Content Questionnaire (JCQ) [53].

In case of an unequal distribution of severe complications (defined as: wound enlargement with more than 8cm or repeat surgery within 2 weeks), between the two study arms, the analyses will be adjusted for these surgery-related characteristics as well.

Cost Measures

The costs will be measured from a societal perspective and consist of: (1) costs of the intervention, (2) health care utilization, and (3) costs associated with lost productivity. All of the costs will be converted to the year 2014 using consumer price indices [54]. The discounting of costs will not be necessary because the follow-up period is limited to one year.

The intervention costs are those that are related to implementing and operating the new care program, and will be estimated using a bottom-up approach. The detailed information regarding the quantity and unit prices of the following resources will be collected: (1) training of involved health care professionals (clinical staff, OP, OT), (2) the eHealth intervention (hosting of webportal, administrator time), and (3) the occupational intervention (number and duration of consultations).

The health care utilization will be assessed on a monthly basis using a retrospective electronic questionnaire. Only the health care costs related to the gynecological surgery will be collected and include: (1) surgery and hospitalization; (2) visit is to health care professionals in primary or secondary care and visits to

alternative medicine therapists; (3) medication; and (4) home care and informal help. If available, Dutch guideline prices will be used to value health care utilization. If cost guidelines are not available, costs will be estimated using real prices or population-based estimates if available in the literature. The prices of the Royal Dutch Society for Pharmacy will be used to value medication [55].

The costs associated with productivity loss consist of absenteeism and presenteeism costs. The absenteeism will be assessed by monthly sick leave calendars. The human capital approach will be used to calculate the costs of losses to production as a result of sick leave due to the gynecologic surgery (net number of days on sick leave during follow-up, multiplied by the estimated prices of production loss of a worker per day of sick leave). The presenteeism (reduced productivity while at work) will be assessed with two items of the Productivity and Disease Questionnaire [56]. A decline in the amount or quality of work performed due to the gynecologic surgery compared to the level at which the patient normally performs, will be considered as presenteeism. The costs associated with presenteeism will be calculated by multiplying the presenteeism score during follow-up by the estimated price of production loss per day.

Process Measures

A process evaluation will be conducted to evaluate the implementation process of the intervention [57]. The assessment of the extent to which the intervention program was applied as intended will provide valuable insight into the facilitators and barriers for future implementation. The process evaluation will take place both on the level of the cluster as well as the patient, and both quantitative and qualitative methods will be used. An automatically generated weblog will enable the analysis of the website activity for each individual participant, giving more insight into which patients used the eHealth intervention, and how it is being used. The appointment system and patient records of the OP will enable us to analyze the number of consultations that have taken place, as well as the reasons for cancellations, and the occurrence of any protocol deviations. By means of an Internet questionnaire at the end of the follow-up

period, patient satisfaction, perceived effectiveness, and any usage barriers will be assessed. The principle investigator will continuously collect reasons for exclusion and dropout during the trial. In accordance to the prior process evaluation conducted [27], the following process measures are included: (1) reach, extent to which the intervention reaches the target population; (2) dose delivered, extent to which the intervention is delivered to the target population; (3) dose received, extent to which the participants used the intervention; (4) fidelity, extent to which the intervention was delivered as planned; and (5) attitudes, satisfaction, perceived effectiveness, and usage barriers.

Cointerventions and Contamination

Cointerventions during the intervention period cannot always be avoided. However, we will be able to determine whether patients received cointerventions by means of the monthly cost diaries. The risk of contamination is reduced by the cluster design of the trial. To assess whether contamination occurred, the patients in both groups are asked about the instructions they received at discharge, which will then be compared to the convalescence recommendations implemented during the intervention phase of the study.

Data Collection

The surgery is considered T_0 . The data will be collected by means of self-reported electronic questionnaires [58] before surgery and 2 weeks (T_1), 6 weeks (T_2), 12 weeks (T_3), 26 weeks (T_4), and 52 weeks (T_5) after surgery. In addition, all of the participants will be requested to fill out a monthly electronic sick leave calendar and cost diary. The patients that are not sick listed, and do not have medical costs during 3 consecutive months, receive a shortened version of the monthly questionnaire. In the case of no response, the patients receive an electronic reminder after 1 and, if necessary, 2 weeks. Every 3 months an attempt will be made to complete missing data regarding RTW, sick leave, and health care usage per email, post, and/or telephone. Table 3 provides an overview of all outcome measures and assessment instruments used in this trial. Not all of the instruments have been validated for Internet use.

Table 3. Assessment of study outcomes.

Outcome measures	–	+	+	+	+	+	
	± 4 weeks	Surgery (T ₀)	2 weeks (T ₁)	6 weeks (T ₂)	3 months (T ₃)	6 months (T ₄)	12 months (T ₅)
Primary							
Duration of sick leave until full sustainable RTW		Monthly sick leave calendar ^a					
Secondary							
Duration of sick leave until first RTW		Monthly sick leave calendar ^a					
Total duration of sick leave		Monthly sick leave calendar ^a					
Recovery (RI-10)	x		x	x	x	x	x
Quality of life (EQ-5D)	x		x	x	x	x	x
Quality of life (SF-36)	x				x	x	x
Self-efficacy (GSES)			x		x		x
Coping (PMS)			x		x		x
Pain intensity (VAS)			x	x	x	x	x
(Post) operative complications			x	x			x ^b
Prognostic factors							
Social demographic variables	x						
Personal factors	x						
Work related factors (DMQ, JCQ)	x						
Type of surgery/complications		x					
Cost							
Care program		Bottom-up approach ^c					
Health care utilization		Monthly cost diary ^a					
Productivity loss		Monthly sick leave calendar ^a					
Process ^d							
Compliance (dose received)		Continuously by weblog					
Attitudes (satisfaction, perceived effectiveness, usage barriers)					x		x
Satisfaction Patient Satisfaction with Occupational Health Services Questionnaire					x		x

^ashort version after 3 consecutive months without sick leave or health care usage

^breview of surgical reports

^ccalculated by research team

^donly intervention group

Blinding

The participants, care providers, and researchers cannot be blinded for the allocated treatment. However, analysis of the data by the researcher will be blind, as all of the patients receive their own study code, under which their data is stored in the database. The assessment of the outcomes is measured through self-reported questionnaires.

Sample Size

We calculated the sample size needed with the method described by Hussey and Hughes [35]. Based on the previous study, we expect a hazard ratio of 1.5 on the primary outcome full sustainable RTW. To achieve a power of 0.8 with a two-tailed alpha of .05, and taking into account a dropout rate of 10%, a

total of 212 patients will be needed when using the log-rank test.

With an intracluster correlation of .05, 9 clusters, and 10 time periods, the design effect is calculated to be 2.14 [35]. By multiplying the design effect by the sample size without a correction for a stepped wedge design, a sample size of 454 women is needed. Assuming that all of the hospitals will include the same amount of participants, each hospital should include approximately 50 patients (5 patients per time period per hospital).

Statistical Analyses

Effect Evaluation

All further described analyses will be performed at the patient level, according to the intention-to-treat principle. In addition, for all tests, a two-tailed significance level of $P \leq .05$ will be considered statistically significant. The statistical software packages that will be used include SPSS (version 16.0) and STATA (version 11.2).

The baseline characteristics will be summarized using descriptive statistics, and compared between the experimental and control group to verify prognostic comparability. In case of coincidental and meaningful differences, these variables will be used as covariates in the further described models.

For the primary outcome, the duration of sick leave until full sustainable RTW, Cox regression analyses will be used to investigate the intervention effect. Both the crude and adjusted analyses will be performed. In the adjusted analyses, the following variables will be used as covariates: (1) hospital, to adjust for clustering (random gamma effect); (2) type of surgery performed; (3) time period, to adjust for naturally occurring changes over time irrespective of the intervention; and (4) optionally, (time period) x (intervention) interaction term, to adjust for time effects (the longer the care program is implemented, the more effective it might be).

The differences in secondary outcomes will be assessed using generalized linear longitudinal mixed models. All of the available measurements (2 weeks, 6 weeks, 12 weeks, 26 weeks, and 52 weeks) will be used, and the baseline scores will be used as covariates, as well as the hospital and the type of surgery (random effect).

To assess whether protocol deviations caused bias, a per protocol analysis will be performed, and the results will be compared to the intention-to-treat analyses. In addition, several subgroup analyses will be performed. The predefined subgroups will be: (1) hysterectomy (TAH, VH, TLH); (2) minimally invasive hysterectomy (VH, TLH); (3) abdominal hysterectomy only; and (4) laparoscopic adnexal surgery only.

Economic Evaluation

Both a cost-effectiveness analysis and a cost-utility analysis will be performed from the societal perspective. The analyses will be performed according to the intention-to-treat principle. The missing cost and effect data will be imputed using multiple imputation [59]. The imputation will include variables that are related to the missing data or the outcome measure, and variables

that differ at baseline between the groups. To account for the skewed distribution of costs, predictive mean matching will be used in the multiple imputation. The number of imputed datasets to be created will be determined based on the fraction of missing information [60]. 'All of the datasets will be analyzed separately, and the results of these analyses will be pooled using Rubin's rules [61]. The incremental cost effectiveness ratios (ICERs) will be calculated by dividing the differences in mean total costs between both treatment groups, by the difference in mean effects between both treatment groups. To avoid double counting, the productivity costs due to sick leave will be excluded in the ICER, with sick leave as the effect measure. The incremental cost utility ratio will be calculated by dividing the incremental costs by the difference in the quality adjusted life years between both treatment groups. To account for the typically skewed distribution of costs, bias corrected and accelerated bootstrapping (5000 replications) will be used to estimate the 95% confidence intervals around the mean cost differences, and the uncertainty surrounding the ICERs. The bootstrapped ICERs will be graphically presented in cost effectiveness planes [62]. The cost effectiveness acceptability curves will be estimated to show the probability of the intervention program to be cost effective in comparison with the usual care for a range of different ceiling ratios, thereby showing decision uncertainty [63]. To assess the robustness of results, several secondary economic analyses will be performed: (1) complete case analysis, (2) per protocol analysis, (3) analysis with costs calculated according to the friction cost approach, and (4) analysis from the health care perspective.

Results

The enrollment of the patients started October 2011. The follow-up period will be completed in August 2014. Data cleaning or analysis has not begun as of this article's submission.

Discussion

Targeting Two Levels

This paper outlines the methodology of a stepped wedge cluster randomized trial to evaluate the cost effectiveness of a care program designed to improve postoperative recovery compared to the usual care. The intervention care program targets two levels: (1) the level of the hospital, and (2) the level of the patient. At the level of the hospital, the newly developed guidelines will be distributed among the clinical staff in order to stimulate evidence-based patient education at the time of discharge. At the patient level, access to an eHealth intervention is provided with tailored convalescence recommendations, and an occupational intervention is available, for those patients at risk of prolonged sick leave, for additional guidance.

What This Study Will Add

The combination of increasing demands on the health care system and the limited health care budget designates a need to enhance the cost effectiveness of our health care system. The introduction of minimally invasive techniques in the last two decades has led to savings in in-hospital care due to shorter lengths of hospital stay, despite higher operative costs, longer

operation time, and more expensive equipment [64-66]. However, early discharge does not necessarily lead to enhanced recovery, as postoperative recovery at home requires a different organization of perioperative care as well, such as preoperative patient education, including the deliverance of evidence-based standardized convalescence recommendations [6,8,9,12,67-70]. As far as we know, our care program is the first intervention developed, and being thoroughly evaluated, that anticipates this transition of perioperative care to the home setting. Second, the utilization of innovative eHealth technologies will limit the workload of involved health care professionals, anticipating a personnel shortage in the health care sector due to a shrinkage of the working population in the near future [71]. Finally, our trial will be one of few that conducted an economic evaluation from a societal perspective, not only taking into account solely direct medical costs—which are important for the hospital perspective—but also including costs associated with postoperative health care utilization and productivity losses due to absenteeism and presenteeism after discharge.

Strengths and Limitations

The main strength of the present study is the choice for a stepped wedge cluster randomized trial. The contamination between study arms is prevented by the cluster design. In addition, the stepped wedge approach enables us to study the implementation process carefully, and gain valuable insight into the facilitators and barriers toward future implementation of the intervention program [72]. Because the crossover of the design is unidirectional, the intervention is not withdrawn from the hospitals during the trial. This is particularly convenient, as our previous trial supports our hypothesis that the care program will lead to enhanced postoperative recovery [73]. Finally, there is a statistical advantage to the stepped wedge approach because the intervention effect is estimated not only by between cluster comparisons, as in a parallel group design, but also by within cluster comparisons, limiting the risk of confounding and increasing statistical power [36,74].

This study also has limitations. First of all, randomized studies without blinding have higher risks of (selection) bias. A second

limitation of this study might be the fact that some of the hospitals have already participated in the earlier trial in 2010. The existing knowledge about the convalescence recommendations could be a source of contamination for the current study, and could lead to an under estimation of the care program effect.

Generalizability

The generalizability of this study will be high, due to the pragmatic study design. In order for procedures to be similar to clinical practice, interference of the research team will be minimized during the trial. The wide diversity of participating (7 teaching, 1 academic, and 1 nonteaching) hospitals, will also contribute to a heterogeneous sample of patients being enrolled in this study, enhancing generalizability. However, we should also be aware of factors that could possibly limit the external validity. A typical feature of eHealth interventions is the risk of selection bias toward the higher educated participants as compared to the general population. Moreover, as the care program was developed in the Dutch setting, and especially tailored to Dutch patients, generalizability of the results of this trial to other countries will be unknown, due to differences in social and health care systems.

Policy Implications

The results of this cost effectiveness study will enable health care policy makers to decide about future implementation of the care program on a broad scale in the Netherlands. In the case that the care program under study is proven to be cost effective, this will have considerable impact. Most importantly, the financial burden on society due to prolonged sick leave following benign gynecological surgery will be substantially reduced. Also, the individual patients will benefit through increased quality of life, and employers will profit because of a decline in absenteeism rates. Moreover, for health care professionals, the care program will be an asset, as it will lead to better organized and more efficient care. Finally, the care program has the potential to maximize the beneficial effects of other recovery enhancing strategies, such as the use of minimally invasive surgery.

Acknowledgments

This study was funded by ZonMw, an organization for health research and development in the Netherlands (project numbers 150020037, 171102015, and 92003590).

Conflicts of Interest

EVB, AVN, HAB, JRA, and JAH are the developers of the care program under study.

Multimedia Appendix 1

Screenshots of ikherstel.

[\[PPT File \(Microsoft PowerPoint Presentation\), 1MB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

CONSORT-EHEALTH checklist V1.6.2 [34].

[\[PDF File \(Adobe PDF File\), 984KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Grant Proposal.

[\[PDF File \(Adobe PDF File\), 150KB-Multimedia Appendix 3\]](#)

Multimedia Appendix 4

Letter of Grant Approval.

[\[PDF File \(Adobe PDF File\), 41KB-Multimedia Appendix 4\]](#)

References

1. Kehlet H, Wilmore DW. Multimodal strategies to improve surgical outcome. *Am J Surg* 2002 Jun;183(6):630-641. [Medline: [12095591](#)]
2. Wodlin NB, Nilsson L. The development of fast-track principles in gynecological surgery. *Acta Obstet Gynecol Scand* 2013 Jan;92(1):17-27. [doi: [10.1111/j.1600-0412.2012.01525.x](#)] [Medline: [22880948](#)]
3. Kjølhed P, Borendal Wodlin N, Nilsson L, Fredrikson M, Wijma K. Impact of stress coping capacity on recovery from abdominal hysterectomy in a fast-track programme: A prospective longitudinal study. *BJOG* 2012 Jul;119(8):998-1006; discussion 1006. [doi: [10.1111/j.1471-0528.2012.03342.x](#)] [Medline: [22568450](#)]
4. Gilmartin J. Contemporary day surgery: Patients' experience of discharge and recovery. *J Clin Nurs* 2007 Jun;16(6):1109-1117. [doi: [10.1111/j.1365-2702.2007.01548.x](#)] [Medline: [17518885](#)]
5. Horvath KJ. Postoperative recovery at home after ambulatory gynecologic laparoscopic surgery. *J Perianesth Nurs* 2003 Oct;18(5):324-334. [Medline: [14569542](#)]
6. Bisgaard T, Klarskov B, Rosenberg J, Kehlet H. Factors determining convalescence after uncomplicated laparoscopic cholecystectomy. *Arch Surg* 2001 Aug;136(8):917-921. [Medline: [11485527](#)]
7. Møller C, Ottesen M, Kehlet H, Ottesen BS. Convalescence recommendations after hysterectomy. A study of opinions among Danish physicians. *Ugeskr Laeger* 2001 Dec 10;163(50):7043-7047. [Medline: [11794035](#)]
8. Naidu M, Sultan AH, Thakar R. Convalescence advice following gynaecological surgery. *J Obstet Gynaecol* 2012 Aug;32(6):556-559. [doi: [10.3109/01443615.2012.693983](#)] [Medline: [22779961](#)]
9. Ottesen M, Møller C, Kehlet H, Ottesen B. Substantial variability in postoperative treatment, and convalescence recommendations following vaginal repair. A nationwide questionnaire study. *Acta Obstet Gynecol Scand* 2001 Nov;80(11):1062-1068. [Medline: [11703209](#)]
10. Vonk Noordegraaf A, Anema JR, Louwse MD, Heymans MW, van Mechelen W, Brölmann HA, et al. Prediction of time to return to work after gynaecological surgery: A prospective cohort study in the Netherlands. *BJOG* 2014 Mar;121(4):487-497. [doi: [10.1111/1471-0528.12494](#)] [Medline: [24245993](#)]
11. Well A, Ijsseldijk A, Bonjer J, Vrijland W, Brouwer K. Medisch Contact. 2002. Boer Kd. Onnodig thuis; ziekteverzuim na liesbreukoperatie URL: <http://medischcontact.artsenet.nl/archief-6/tijdschriftartikel/05311/onnodig-thuis.htm> [accessed 2014-06-05] [WebCite Cache ID [6Q6M43leM](#)]
12. Carter E. Ready for home? *Nurs Times* 1981 May 7;77(19):826-829. [Medline: [6909791](#)]
13. Evenson M, Payne D, Nygaard I. Recovery at home after major gynecologic surgery: How do our patients fare? *Obstet Gynecol* 2012 Apr;119(4):780-784. [doi: [10.1097/AOG.0b013e31824bb15e](#)] [Medline: [22433341](#)]
14. Majeed AW, Brown S, Williams N, Hannay DR, Johnson AG. Variations in medical attitudes to postoperative recovery period. *BMJ* 1995 Jul 29;311(7000):296 [FREE Full text] [Medline: [7633239](#)]
15. Easton K, Read MD, Woodman NM. Influence of early discharge after hysterectomy on patient outcome and GP workloads. *J Obstet Gynaecol* 2003 May;23(3):271-275. [Medline: [12850859](#)]
16. Johansen P, Al-Khafagi SK, Thøstesen LM, Lauszus FF, Rasmussen KL. Analysis of need for sick leave after hysterectomy. *Ugeskr Laeger* 2008 Apr 21;170(17):1465-1468. [Medline: [18462627](#)]
17. Rasmussen KL, Hansen V, Madzak F, Ljungstrøm B, Lauszus FF. Feeling of illness after hysterectomy. Women's own assessment. *Ugeskr Laeger* 2001 Dec 10;163(50):7040-7042. [Medline: [11794034](#)]
18. van Oostrom SH, Driessen MT, de Vet HC, Franche RL, Schonstein E, Loisel P, et al. Workplace interventions for preventing work disability. *Cochrane Database Syst Rev* 2009(2):CD006955. [doi: [10.1002/14651858.CD006955.pub2](#)] [Medline: [19370664](#)]
19. Waddell G, Burton AK. Is work good for your health and well-being?. 2006 URL: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/214326/hwwb-is-work-good-for-you.pdf [accessed 2014-03-03] [WebCite Cache ID [6NnRlwNCh](#)]
20. Reeuwijk KG, Robroek SJ, Hakkaart L, Burdorf A. How work impairments and reduced work ability are associated with health care use in workers with musculoskeletal disorders, cardiovascular disorders or mental disorders. *J Occup Rehabil* 2014 Jan 4. [doi: [10.1007/s10926-013-9492-3](#)] [Medline: [24390780](#)]
21. Statline Databank. Surgeries in the Netherlands. 2014 February URL: <http://www.webcitation.org/6Nm4SznjS>

22. Vonk Noordegraaf A, Huirne JA, Brölmann HA, van Mechelen W, Anema JR. Multidisciplinary convalescence recommendations after gynaecological surgery: A modified Delphi method among experts. *BJOG* 2011 Dec;118(13):1557-1567. [doi: [10.1111/j.1471-0528.2011.03091.x](https://doi.org/10.1111/j.1471-0528.2011.03091.x)] [Medline: [21895950](https://pubmed.ncbi.nlm.nih.gov/21895950/)]
23. Vonk Noordegraaf A, Huirne JA, Pittens CA, van Mechelen W, Broerse JE, Brölmann HA, et al. eHealth program to empower patients in returning to normal activities and work after gynecological surgery: Intervention mapping as a useful method for development. *J Med Internet Res* 2012;14(5):e124 [FREE Full text] [doi: [10.2196/jmir.1915](https://doi.org/10.2196/jmir.1915)] [Medline: [23086834](https://pubmed.ncbi.nlm.nih.gov/23086834/)]
24. Pittens CA, Vonk Noordegraaf A, van Veen SC, Anema JR, Huirne JA, Broerse JE. The involvement of gynaecological patients in the development of a clinical guideline for resumption of (work) activities in the Netherlands. *Health Expect* 2013 Aug 29. [doi: [10.1111/hex.12121](https://doi.org/10.1111/hex.12121)] [Medline: [23992108](https://pubmed.ncbi.nlm.nih.gov/23992108/)]
25. Vonk Noordegraaf A, Huirne JA, Brölmann HA, Emanuel MH, van Kesteren PJ, Kleiverda G, et al. Effectiveness of a multidisciplinary care program on recovery and return to work of patients after gynaecological surgery; design of a randomized controlled trial. *BMC Health Serv Res* 2012;12:29 [FREE Full text] [doi: [10.1186/1472-6963-12-29](https://doi.org/10.1186/1472-6963-12-29)] [Medline: [22296950](https://pubmed.ncbi.nlm.nih.gov/22296950/)]
26. Vonk Noordegraaf A, Anema J, van Mechelen W, Knol D, van Baal W, van Kesteren P, et al. A personalised eHealth programme reduces the duration until return to work after gynaecological surgery: Results of a multicentre randomised trial. *BJOG* 2014 Feb 11. [doi: [10.1111/1471-0528.12661](https://doi.org/10.1111/1471-0528.12661)] [Medline: [24511914](https://pubmed.ncbi.nlm.nih.gov/24511914/)]
27. Bouwsma EV, Vonk Noordegraaf A, Szlávik Z, Brölmann HA, Emanuel MH, Lips JP, et al. Process evaluation of a multidisciplinary care program for patients undergoing gynaecological surgery. *J Occup Rehabil* 2013 Sep 22. [doi: [10.1007/s10926-013-9475-4](https://doi.org/10.1007/s10926-013-9475-4)] [Medline: [24057871](https://pubmed.ncbi.nlm.nih.gov/24057871/)]
28. Drummond MF, Sculpher MJ, Torrance GW, O'Brien BJ, Stoddart GL. *Methods for the economic evaluation of health care programmes*. Oxford: Oxford University Press; 2005.
29. Chan AW, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin JA, et al. SPIRIT 2013 explanation and elaboration: Guidance for protocols of clinical trials. *BMJ* 2013;346:e7586 [FREE Full text] [Medline: [23303884](https://pubmed.ncbi.nlm.nih.gov/23303884/)]
30. Schulz KF, Altman DG, Moher D, CONSORT Group. CONSORT 2010 statement: Updated guidelines for reporting parallel group randomized trials. *Obstet Gynecol* 2010 May;115(5):1063-1070. [doi: [10.1097/AOG.0b013e3181d9d421](https://doi.org/10.1097/AOG.0b013e3181d9d421)] [Medline: [20410783](https://pubmed.ncbi.nlm.nih.gov/20410783/)]
31. Moher D, Hopewell S, Schulz KF, Montori V, Gøtzsche PC, Devereaux PJ, et al. CONSORT 2010 explanation and elaboration: Updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;340:c869 [FREE Full text] [Medline: [20332511](https://pubmed.ncbi.nlm.nih.gov/20332511/)]
32. Campbell MK, Piaggio G, Elbourne DR, Altman DG, CONSORT Group. Consort 2010 statement: Extension to cluster randomised trials. *BMJ* 2012;345:e5661. [Medline: [22951546](https://pubmed.ncbi.nlm.nih.gov/22951546/)]
33. Eysenbach G, CONSORT-EHEALTH Group. CONSORT-EHEALTH: Improving and standardizing evaluation reports of Web-based and mobile health interventions. *J Med Internet Res* 2011;13(4):e126 [FREE Full text] [doi: [10.2196/jmir.1923](https://doi.org/10.2196/jmir.1923)] [Medline: [22209829](https://pubmed.ncbi.nlm.nih.gov/22209829/)]
34. Brown CA, Lilford RJ. The stepped wedge trial design: A systematic review. *BMC Med Res Methodol* 2006;6:54 [FREE Full text] [doi: [10.1186/1471-2288-6-54](https://doi.org/10.1186/1471-2288-6-54)] [Medline: [17092344](https://pubmed.ncbi.nlm.nih.gov/17092344/)]
35. Hussey MA, Hughes JP. Design and analysis of stepped wedge cluster randomized trials. *Contemp Clin Trials* 2007 Feb;28(2):182-191. [doi: [10.1016/j.cct.2006.05.007](https://doi.org/10.1016/j.cct.2006.05.007)] [Medline: [16829207](https://pubmed.ncbi.nlm.nih.gov/16829207/)]
36. Mdege ND, Man MS, Taylor Nee Brown CA, Torgerson DJ. Systematic review of stepped wedge cluster randomized trials shows that design is particularly used to evaluate interventions during routine implementation. *J Clin Epidemiol* 2011 Sep;64(9):936-948. [doi: [10.1016/j.jclinepi.2010.12.003](https://doi.org/10.1016/j.jclinepi.2010.12.003)] [Medline: [21411284](https://pubmed.ncbi.nlm.nih.gov/21411284/)]
37. de Vries H, Dijkstra M, Kuhlman P. Self-efficacy: The third factor besides attitude and subjective norm as a predictor of behavioural intentions. *Health Educ Res* 1988;3(3):273-282. [doi: [10.1093/her/3.3.273](https://doi.org/10.1093/her/3.3.273)]
38. Lechner L, Kremers S, Meertens R, de Vries H. Determinanten van gedrag. In: *Gezondheidsvoorlichting en gedragsverandering Health Education and Behavioral Change*. 8 ed. Assen: van Gorcum; 2012:75-106.
39. ikherstel. Welkom bij ikherstel.nl URL: <http://www.ikherstel.nl/> [accessed 2014-06-04] [WebCite Cache ID [6Q5N3xli7](https://www.webcitation.org/6Q5N3xli7)]
40. van Oostrom SH, van Mechelen W, Terluin B, de Vet HC, Knol DL, Anema JR. A workplace intervention for sick-listed employees with distress: Results of a randomised controlled trial. *Occup Environ Med* 2010 Sep;67(9):596-602. [doi: [10.1136/oem.2009.050849](https://doi.org/10.1136/oem.2009.050849)] [Medline: [20363809](https://pubmed.ncbi.nlm.nih.gov/20363809/)]
41. Vermeulen SJ, Anema JR, Schellart AJ, Knol DL, van Mechelen W, van der Beek AJ. A participatory return-to-work intervention for temporary agency workers and unemployed workers sick-listed due to musculoskeletal disorders: Results of a randomized controlled trial. *J Occup Rehabil* 2011 Sep;21(3):313-324 [FREE Full text] [doi: [10.1007/s10926-011-9291-7](https://doi.org/10.1007/s10926-011-9291-7)] [Medline: [21336673](https://pubmed.ncbi.nlm.nih.gov/21336673/)]
42. Anema JR, Steenstra IA, Urlings IJ, Bongers PM, de Vroome EM, van Mechelen W. Participatory ergonomics as a return-to-work intervention: A future challenge? *Am J Ind Med* 2003 Sep;44(3):273-281. [doi: [10.1002/ajim.10259](https://doi.org/10.1002/ajim.10259)] [Medline: [12929147](https://pubmed.ncbi.nlm.nih.gov/12929147/)]
43. de Jong AM, Vink P. Participatory ergonomics applied in installation work. *Appl Ergon* 2002 Sep;33(5):439-448. [Medline: [12236653](https://pubmed.ncbi.nlm.nih.gov/12236653/)]

44. de Vet HC, Heymans MW, Dunn KM, Pope DP, van der Beek AJ, Macfarlane GJ, et al. Episodes of low back pain: A proposal for uniform definitions to be used in research. *Spine (Phila Pa 1976)* 2002 Nov 1;27(21):2409-2416. [doi: [10.1097/01.BRS.0000030307.34002.BE](https://doi.org/10.1097/01.BRS.0000030307.34002.BE)] [Medline: [12438991](https://pubmed.ncbi.nlm.nih.gov/12438991/)]
45. Kluivers KB, Hendriks JC, Mol BW, Bongers MY, Vierhout ME, Brölmann HA, et al. Clinimetric properties of 3 instruments measuring postoperative recovery in a gynecologic surgical population. *Surgery* 2008 Jul;144(1):12-21. [doi: [10.1016/j.surg.2008.03.027](https://doi.org/10.1016/j.surg.2008.03.027)] [Medline: [18571580](https://pubmed.ncbi.nlm.nih.gov/18571580/)]
46. Dolan P. Modeling valuations for EuroQol health states. *Med Care* 1997 Nov;35(11):1095-1108. [Medline: [9366889](https://pubmed.ncbi.nlm.nih.gov/9366889/)]
47. Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderman R, et al. Translation, validation, and norming of the Dutch language version of the SF-36 health survey in community and chronic disease populations. *J Clin Epidemiol* 1998 Nov;51(11):1055-1068. [Medline: [9817123](https://pubmed.ncbi.nlm.nih.gov/9817123/)]
48. Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992 Jun;30(6):473-483. [Medline: [1593914](https://pubmed.ncbi.nlm.nih.gov/1593914/)]
49. Teeuw B, Schwarzer R, Jerusalem M. Dutch adaptation of the general self-efficacy scale. 1994 URL: http://userpage.fu-berlin.de/~gesund/publicat/ehps_cd/health/dutch.htm [accessed 2014-03-03] [WebCite Cache ID [6NnSKKgm3](https://www.webcitation.org/6NnSKKgm3)]
50. Pearlin LI, Schooler C. The structure of coping. *J Health Soc Behav* 1978 Mar;19(1):2-21. [Medline: [649936](https://pubmed.ncbi.nlm.nih.gov/649936/)]
51. Von Korff M, Ormel J, Keefe FJ, Dworkin SF. Grading the severity of chronic pain. *Pain* 1992 Aug;50(2):133-149. [Medline: [1408309](https://pubmed.ncbi.nlm.nih.gov/1408309/)]
52. Hildebrandt VH, Bongers PM, van Dijk FJ, Kemper HC, Dul J. Dutch musculoskeletal questionnaire: Description and basic qualities. *Ergonomics* 2001 Oct 10;44(12):1038-1055. [doi: [10.1080/00140130110087437](https://doi.org/10.1080/00140130110087437)] [Medline: [11780727](https://pubmed.ncbi.nlm.nih.gov/11780727/)]
53. Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. The job content questionnaire (JCQ): An instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol* 1998 Oct;3(4):322-355. [Medline: [9805280](https://pubmed.ncbi.nlm.nih.gov/9805280/)]
54. Statline Databank. Consumer prices. URL: <http://www.cbs.nl/nl-NL/menu/themas/inkomen-bestedingen/cijfers/default.htm> [accessed 2014-03-02] [WebCite Cache ID [6NluSnQ0M](https://www.webcitation.org/6NluSnQ0M)]
55. Z-index. G-standaard. URL: <http://www.z-index.nl/g-standaard/g-standaard> [accessed 2014-03-03] [WebCite Cache ID [6NnSqBHJW](https://www.webcitation.org/6NnSqBHJW)]
56. Koopmanschap MA. PRODISQ: A modular questionnaire on productivity and disease for economic evaluation studies. *Expert Rev Pharmacoecon Outcomes Res* 2005 Feb;5(1):23-28. [doi: [10.1586/14737167.5.1.23](https://doi.org/10.1586/14737167.5.1.23)] [Medline: [19807557](https://pubmed.ncbi.nlm.nih.gov/19807557/)]
57. Steckler AB, Linnan L. Process evaluation for public health interventions and research: An overview. In: *Process evaluation for public health interventions and research*. USA: Jossey-Bass; 2002.
58. Net Q. Utrecht, The Netherlands: NetQuestionnaires Nederland BV; 2011. Software for creating assessment of internet surveys. Computer program URL: <http://www.netq-enquete.nl/nl> [accessed 2014-06-07] [WebCite Cache ID [6QA6xaD7e](https://www.webcitation.org/6QA6xaD7e)]
59. van Buuren S, Boshuizen HC, Knook DL. Multiple imputation of missing blood pressure covariates in survival analysis. *Stat Med* 1999 Mar 30;18(6):681-694. [Medline: [10204197](https://pubmed.ncbi.nlm.nih.gov/10204197/)]
60. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011 Feb 20;30(4):377-399. [doi: [10.1002/sim.4067](https://doi.org/10.1002/sim.4067)] [Medline: [21225900](https://pubmed.ncbi.nlm.nih.gov/21225900/)]
61. Rubin DB. *Multiple imputation for nonresponse in surveys*. New York: Wiley; 1987.
62. Black WC. The CE plane: A graphic representation of cost-effectiveness. *Med Decis Making* 1990;10(3):212-214. [Medline: [2115096](https://pubmed.ncbi.nlm.nih.gov/2115096/)]
63. Fenwick E, O'Brien BJ, Briggs A. Cost-effectiveness acceptability curves--facts, fallacies and frequently asked questions. *Health Econ* 2004 May;13(5):405-415. [doi: [10.1002/hec.903](https://doi.org/10.1002/hec.903)] [Medline: [15127421](https://pubmed.ncbi.nlm.nih.gov/15127421/)]
64. Bijen CB, Vermeulen KM, Mourits MJ, de Bock GH. Costs and effects of abdominal versus laparoscopic hysterectomy: Systematic review of controlled trials. *PLoS One* 2009;4(10):e7340 [FREE Full text] [doi: [10.1371/journal.pone.0007340](https://doi.org/10.1371/journal.pone.0007340)] [Medline: [19806210](https://pubmed.ncbi.nlm.nih.gov/19806210/)]
65. Nieboer TE, Johnson N, Lethaby A, Tavender E, Curr E, Garry R, et al. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev* 2009(3):CD003677. [doi: [10.1002/14651858.CD003677.pub4](https://doi.org/10.1002/14651858.CD003677.pub4)] [Medline: [19588344](https://pubmed.ncbi.nlm.nih.gov/19588344/)]
66. Sculpher M, Manca A, Abbott J, Fountain J, Mason S, Garry R. Cost effectiveness analysis of laparoscopic hysterectomy compared with standard hysterectomy: Results from a randomised trial. *BMJ* 2004 Jan 17;328(7432):134 [FREE Full text] [doi: [10.1136/bmj.37942.601331.EE](https://doi.org/10.1136/bmj.37942.601331.EE)] [Medline: [14711748](https://pubmed.ncbi.nlm.nih.gov/14711748/)]
67. Bay-Nielsen M, Bisgaard T. Convalescence and sick leave following inguinal hernia repair. *Ugeskr Laeger* 2009 Sep 28;171(40):2899-2901. [Medline: [19814936](https://pubmed.ncbi.nlm.nih.gov/19814936/)]
68. Callesen T, Klarskov B, Bech K, Kehlet H. Short convalescence after inguinal herniorrhaphy with standardised recommendations: Duration and reasons for delayed return to work. *Eur J Surg* 1999 Mar;165(3):236-241. [doi: [10.1080/110241599750007108](https://doi.org/10.1080/110241599750007108)] [Medline: [10231657](https://pubmed.ncbi.nlm.nih.gov/10231657/)]
69. Clayton M, Verow P. Advice given to patients about return to work and driving following surgery. *Occup Med (Lond)* 2007 Oct;57(7):488-491 [FREE Full text] [doi: [10.1093/occmed/kqm063](https://doi.org/10.1093/occmed/kqm063)] [Medline: [17906266](https://pubmed.ncbi.nlm.nih.gov/17906266/)]

70. Berendsen AJ, de Jong GM, Meyboom-de Jong B, Dekker JH, Schuling J. Transition of care: Experiences and preferences of patients across the primary/secondary interface - a qualitative study. *BMC Health Serv Res* 2009;9:62 [FREE Full text] [doi: [10.1186/1472-6963-9-62](https://doi.org/10.1186/1472-6963-9-62)] [Medline: [19351407](https://pubmed.ncbi.nlm.nih.gov/19351407/)]
71. Noteboom A, Blankers I, Goudriaan R, Groot W. E-health en zelfmanagement: een panacee voor arbeidstekorten en kostenoverschrijdingen in de zorg?. URL: <http://www.ape.nl/include/downloadFile.asp?id=286> [accessed 2014-03-03] [WebCite Cache ID 6NnNevrIF]
72. Dekkers OM. The stepped wedge design. *Ned Tijdschr Geneesk* 2012;156(9):A4069. [Medline: [22373553](https://pubmed.ncbi.nlm.nih.gov/22373553/)]
73. Edwards SJ, Braunholtz DA, Lilford RJ, Stevens AJ. Ethical issues in the design and conduct of cluster randomised controlled trials. *BMJ* 1999 May 22;318(7195):1407-1409 [FREE Full text] [Medline: [10334756](https://pubmed.ncbi.nlm.nih.gov/10334756/)]
74. Woertman W, de Hoop E, Moerbeek M, Zuidema SU, Gerritsen DL, Teerenstra S. Stepped wedge designs could reduce the required sample size in cluster randomized trials. *J Clin Epidemiol* 2013 Jul;66(7):752-758. [doi: [10.1016/j.jclinepi.2013.01.009](https://doi.org/10.1016/j.jclinepi.2013.01.009)] [Medline: [23523551](https://pubmed.ncbi.nlm.nih.gov/23523551/)]

Abbreviations

CONSORT: CONSolidated Standards Of Reporting Trials

DMQ: Dutch Musculoskeletal Questionnaire

EQ-5D: EuroQol-5D

GP: general physician

GSES: General Self-Efficacy Scale

ICER: incremental cost effectiveness ratio

IQR: interquartile range

JCQ: Job Content Questionnaire

LAS: laparoscopic adnexal surgery

OP: occupational physician

OT: occupational therapist

PMS: Pearlin Mastery Scale

RI-10: Recovery Index-10

RTW: return to work

SF-36: Short-Form Health Survey

TAH: total abdominal hysterectomy

TLH: total laparoscopic hysterectomy

TP: time period

VAS: Visual Analogue Scale

VH: vaginal hysterectomy

Edited by G Eysenbach; submitted 01.04.14; peer-reviewed by G Kok; comments to author 22.04.14; accepted 21.05.14; published 18.06.14

Please cite as:

Bouwsma EVA, Anema JR, Vonk Noordegraaf A, Knol DL, Bosmans JE, Schraffordt Koops SE, van Kesteren PJM, van Baal WM, Lips JP, Emanuel MH, Scholten PC, Mozes A, Adriaanse AH, Brölmann HAM, Huirne JAF

The Cost Effectiveness of a Tailored, Web-Based Care Program to Enhance Postoperative Recovery in Gynecologic Patients in Comparison With Usual Care: Protocol of a Stepped Wedge Cluster Randomized Controlled Trial

JMIR Res Protoc 2014;3(2):e30

URL: <http://www.researchprotocols.org/2014/2/e30/>

doi: [10.2196/resprot.3236](https://doi.org/10.2196/resprot.3236)

PMID: [24943277](https://pubmed.ncbi.nlm.nih.gov/24943277/)

©Esther VA Bouwsma, Johannes R Anema, Antonie Vonk Noordegraaf, Dirk L Knol, Judith E Bosmans, Steven E Schraffordt Koops, Paul JM van Kesteren, W Marchien van Baal, Jos P Lips, Mark H Emanuel, Petrus C Scholten, Alexander Mozes, Albert H Adriaanse, Hans AM Brölmann, Judith AF Huirne. Originally published in *JMIR Research Protocols* (<http://www.researchprotocols.org>), 18.06.2014. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Research Protocols*, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.