

Protocol

# Mobile Phone Apps for University Students With Hazardous Alcohol Use: Study Protocol for Two Consecutive Randomized Controlled Trials

Anne H Berman<sup>1</sup>, PhD; Mikael Gajecki<sup>1</sup>, MSc; Morgan Fredriksson<sup>2</sup>; Kristina Sinadinovic<sup>1</sup>, PhD; Claes Andersson<sup>3</sup>, PhD

<sup>1</sup>Center for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden

<sup>2</sup>Liquid Media AB, Stockholm, Sweden

<sup>3</sup>Department of Criminology, Malmö University, Malmö, Sweden

**Corresponding Author:**

Anne H Berman, PhD

Center for Psychiatry Research

Department of Clinical Neuroscience

Karolinska Institutet

Norra Stationsgatan 69, plan 7

Stockholm, 11364

Sweden

Phone: 46 704245360

Fax: 46 8311101

Email: [anne.h.berman@ki.se](mailto:anne.h.berman@ki.se)

## Abstract

**Background:** About 50% of university students overconsume alcohol, and drinking habits in later adulthood are to some extent established during higher educational studies. Several studies have demonstrated that Internet-based interventions have positive effects on drinking habits among university students. Our recent study evaluated two mobile phone apps targeting drinking choices at party occasions via personalized feedback on estimated blood alcohol concentration (eBAC) for students with hazardous drinking. No changes in drinking parameters were found over a seven-week period apart from an increase in number of drinking occasions among men for one of the apps tested. Up to 30% of the study participants drank at potentially harmful levels: higher than the national recommended number of standard drinks per week (a maximum of 9 for women and 14 for men) in Sweden.

**Objective:** (1) To evaluate improved versions of the two mobile phone apps tested in our prior trial, in a new, 3-armed randomized controlled trial among university students with at least hazardous drinking habits according to the Alcohol Use Disorders Identifications Test (AUDIT; Study 1). (2) After 6 weeks, to target study participants showing alcohol consumption higher than the national recommended levels for standard drinks per week by offering them participation in a second, 2-armed randomized trial evaluating an additional mobile phone app with skill enhancement tasks (Study 2). (3) To follow participants at 6, 12 and 18 weeks after recruitment to Study 1 and at 6 and 12 weeks after recruitment to Study 2.

**Methods:** Two randomized controlled trials are conducted. Study 1: Students are recruited at four Swedish universities, via direct e-mail and advertisements on Facebook and student union web sites. Those who provide informed consent, have a mobile phone, and show at least hazardous alcohol consumption according to the AUDIT ( $\geq 6$  for women;  $\geq 8$  points for men) are randomized into three groups. Group 1 has access to the Swedish government alcohol monopoly's app, Promillekoll, offering real-time estimated eBAC calculation; Group 2 has access to a Web-based app, PartyPlanner, developed by the research group, offering real-time eBAC calculation with planning and follow-up functions; and Group 3 participants are controls. Follow-up is conducted at 6, 12 and 18 weeks. Study 2. Participants who at the first 6-week follow-up show drinking levels higher than 9 (W) or 14 (M) standard drinks (12 g alcohol) per week, are offered participation in Study 2. Those who consent are randomized to either access to a skills training app, TeleCoach or to a wait-list control group.

**Results:** Latent Markov models for Study 1 and mixed models analyses for Study 2 will be performed. Study 2 data will be analyzed for publication during the spring of 2016; Study 1 data will be analyzed for publication during the fall of 2016.

**Conclusions:** If mobile phone interventions for reducing hazardous alcohol use are found to be effective, the prospects for positively influencing substance use-related health among university students can considerably improve.

**Trial Registration:** ClinicalTrials.gov <http://clinicaltrials.gov/ct2/show/NCT02064998> (Archived by WebCite at <http://www.webcitation.org/6dy0AIVRP>)

(*JMIR Res Protoc* 2015;4(4):e139) doi: [10.2196/resprot.4894](https://doi.org/10.2196/resprot.4894)

## KEYWORDS

randomized controlled trial, universities, alcohol abuse, prevention, mobile phone, eHealth, mHealth

## Introduction

Research studies have shown that consumption of alcohol is higher during the years at university than at any other age [1]. Furthermore, it appears that students at college and university establish their future adult alcohol habits [2]. Individual, social and environmental factors affect this development [3]. Several studies have demonstrated that effective intervention approaches, both at individual and group levels, are available to reduce hazardous drinking among students. These methods are well documented internationally [4,5] and as well as nationally in a meta-analysis [6].

A major behavior change component for reducing problematic alcohol use in general is screening with brief intervention (SBI), where the primary goal is to moderate consumption to sensible levels and to eliminate harmful drinking practices in order to reduce negative outcomes of drinking [7]. At minimum, SBI includes only screening of drinking habits, followed by clear and direct feedback. It can also include detailed conversations covering setting up goals, use of behavioral modification techniques, self-help exercises, and continual reinforcement [8]. For university students, effective interventions are termed brief motivational interventions (BMIs), providing personalized feedback on individual drinking habits and their consequences, based on self-monitoring, as well as exploration of motives for using alcohol. Specific behavior change components include the personalized feedback aspect, particularly normative feedback, in relation to students in the same university context; studies of feedback on Blood Alcohol Concentration (BAC) have shown mixed support [9].

## New Technology

Despite availability of interventions, only a small percentage of students seek help for problem drinking [10]. College students with heavy episodic drinking (HED) at least once a month have been found to prefer computerized methods [11]. Digital interventions for alcohol problems, regardless of delivery mode, offer small but meaningful effects [12-18]. They are as effective as alternative interventions offered face-to-face by a live counselor when compared with controls but, in direct comparisons, face-to-face interventions have been shown to be more effective [17,18].

## Mobile Apps

In recent years, mobile phones have offered constant access to hand held computers. Easy to use mobile phone apps can fill a variety of functions, such as games and other entertainment. Depending on its function, the app may or may not require an Internet connection. Apps have been used for registering weight for obesity control [19] and, in a guided version, to help users

with behavioral activation in their own chosen valued direction [20]. Research on mobile phone apps for reducing alcohol consumption is in its infancy. Over 3000 apps on alcohol-related topics are available, but these usually have no therapeutic purpose and even provide incorrect information, for example regarding blood alcohol content [21]. An analysis of user experiences of 87 Android apps showed, however, that these in some cases inspire users to keep their alcohol use down [22]. A very recent content analysis of 800 alcohol apps with a focus on behavior change techniques found that the majority “implicitly or explicitly promoted the use of alcohol”. Of 61 apps coded for behavior change techniques, the most frequent techniques used (over 40%) were self-monitoring, information on negative consequences of alcohol use and positive consequences of abstinence, and personalized feedback. The analysis cites no research specifically evaluating the apps reviewed [23].

Indeed, systematic research on mobile phone apps for university students has so far been very sparse [24]. One study targeted smoking and HED based on BASICS (Brief Alcohol Screening and Intervention for College Students) and found no reduction over a one-month period in HED, although students who completed more intervention modules were less likely to drink during the initial 14-day assessment period in the study [25]. In a second study, our own research group conducted a 3-armed randomized controlled trial among university students with problematic drinking, comparing the effects of 2 different apps, one offering personalized feedback on estimated blood alcohol concentration (eBAC) at live drinking occasions and the other offering planning, live monitoring of eBAC with personalized feedback, and follow-up of specific drinking occasions based on eBAC (see [Figure 1](#) for an overview of app components). We found no changes in drinking parameters over a 7-week period apart from an *increase* in number of drinking occasions among men for one of the apps tested. Also, dropping out from the study was associated with drinking at levels higher than the recommended maximum number of standard drinks per week (9 for women and 14 for men); students who drank at higher than recommended levels comprised a sub-group of almost one-third of students assessed as having at least hazardous drinking habits according to the Alcohol Use Disorders Identification Test (AUDIT) [26].

In recent years, alcohol apps for mobile phones have seen an exponential rise in growth. Their potential is high in view of their extensive reach in the population, but the evidence for their effectiveness in reducing problematic alcohol use is lacking [27]. Questions that remain to be answered are whether it is possible to reduce hazardous or even harmful drinking with apps, as well what components an app actually needs to include in order to be effective.

**Figure 1.** Behavior change components included in the two apps used in Study 1 and in the app used in Study 2.

Behavior Change Components	Promillekoll	PartyPlanner	TeleCoach
Providing information on Protective Behavioral Strategies (PBS)	X	X	X
Planning future parties		X	
Realtime feedback on blood alcohol count (BAC)	X	X	
Comparative follow-up ("How did it go?")		X	
Registering intake over time		(X)	X
Skill training for saying no to alcohol and feeling better without alcohol			X

## Study Aim

This research has two principal aims. First, we aim to evaluate mobile phone apps in improved, revised versions compared to our previous study [26] for university students with problematic levels of alcohol use based on AUDIT scores for at least hazardous drinking. We also aim to extend follow-up beyond 6 weeks, to 12 and 18 weeks. We hypothesize that the improved app versions may show positive effects with reduced alcohol use over time in comparison to an untreated control group, in contrast to no effects in the 6-week follow-up in our previous study [26]. Secondly, we aim to target the needs of a sub-group of university students with particularly high alcohol consumption according to national public health thresholds for unhealthy drinking, by evaluating a new mobile phone app with skills enhancement functions. We hypothesize that dropout over time will be reduced for students in this sub-group, who are offered an app addressing their specific, more extensive need of help to control drinking. Also, we hypothesize that students offered this app will reduce their drinking to a larger extent than students in a wait-list control group as well as those in an untreated

control group (separately culled from the wider data set for Study 1). These aims will be addressed through two consecutive randomized trials (Trial Registration at ClinicalTrials.gov, identifier NCT02064998).

The specific research questions are as follows:

**Study 1:** Among university students with at least hazardous alcohol use according to the AUDIT, does access and self-reported use of two different mobile phone apps (see Figure 2) lead to reduced alcohol consumption in comparison to an untreated, assessment only control group, at 6, 12 and 18 weeks after registration for the study?

**Study 2:** Among university students who drink over the recommended 9 and 14 standard drinks per week for women and men, respectively, does access and self-reported use of the TeleCoach app (see Figure 2) lead to lower proportions of individuals with consumption levels under the recommended number of drinks per week, in comparison to a wait-list control group and an untreated control group, 6 and 12 weeks after registration to Study 2 (12 and 18 weeks after registration for Study 1)?



risk of harmful consequences, and is of particular concern since 1 out of 4 deaths among individuals 15-29 years old in the EU are alcohol-related [31]. These students are given feedback on their elevated consumption level with higher risk of harmful consequences. They continue to have access to the app they were randomized to in Study 1, but are randomized to one of two groups: access to an in-depth self-help TeleCoach app that the research group has developed, or to a wait-list control group who will be given access to the app at the 12-week follow-up.

All study participants who complete the final follow-up will participate in a lottery of 3 iPad devices, through a partnership with Save the Children.

## Interventions

All three app interventions used in Studies 1 and/or 2 are described below. Figure 1 shows a simplified overview of basic behavior change components included in each app.

### *Promillekoll App (Translation: “Check your BAC”)*

This mobile phone app is available in iPhone and Android versions, and allows the user to register alcohol consumption in real time, giving immediate feedback on the eBAC. The app alerts the user if s/he surpasses an alcohol concentration of 0.06 percent BAC, a level where negative consequences can begin to occur, and only displays values up to 0.08 percent, in an effort to emphasize the message to users that higher eBAC levels are harmful. The app also provides separate information texts on alcohol and BAC. The app was publicly launched by the Swedish Alcohol Monopoly (Systembolaget) in the autumn of 2012 and improved a year later on the basis of the Monopoly's own surveys and 343 student comments from our study in the spring of 2013 (unpublished data). The Promillekoll app is theoretically based on the assumption that information about one's own real-time eBAC levels can contribute to protective cognitive and behavioral strategies. A further mechanism, congruent with the Theory of Planned Behavior (TPB) [32], is that providing information and feedback on risky levels of eBAC modifies the intention to consume alcohol. Promillekoll also offers selected specific protective behavioral strategies (PBS) [33] to maintain alcohol consumption at or below the 0.06 percent BAC level. No user data are collected.

### *PartyPlanner App*

This Web-based app guides the user to plan drinking occasions so the consumption level stays below risky levels of eBAC, at about 0.06 percent. The app also allows the user to record real-time consumption and provides immediate feedback on

how to adjust their consumption to maintain a healthy eBAC level to 0.06 percent or less. If the user has created a plan for a specific drinking occasion and then followed it up, the app offers a graphic visual comparison of the plan, with the logged real-time event after the actual drinking occasion. These app components are in line with the goal setting and personalized feedback components of successful BMIs [9].


The app was built by the research team in the fall of 2012 and revised in the fall of 2014 for the current study. In the revised version, users receive feedback about the health benefits of lower alcohol use and potential social losses connected to over-consumption, since laboratory studies have shown that this kind of information affects men's drinking intentions in a downward direction [34]; we focused on adapting information primarily to men because we saw a significant negative effect for men using Promillekoll in our first study, and unpublished secondary analyses suggested that men might have more difficulty using an app constructively in a social context where alcohol is consumed, than women. An example of the type of feedback given is shown in Figure 3. PartyPlanner also includes several PBS [33] to maintain alcohol consumption at a minimally harmful level.

Note: The level for non-risky eBAC is below 0.06 percent for both Promillekoll and PartyPlanner, in correlation with guidelines used in the BASICS program [35]; this level is somewhat more conservative than the 0.08 percent level correlated with the concept of binge drinking [36].

### *TeleCoach App*

This app is based on an Interactive Voice Response (IVR) automated telephony intervention developed by the research group in 2009-2012 [37]. Aside from offering skill acquisition for PBS [33], it provides skill training exercises for individuals who want to reduce or end their alcohol consumption but have experienced trouble in achieving these behavioral changes. The app includes a behavioral chart, and modules on “saying no to alcohol” and “feeling better without alcohol.” The “saying no” module includes instruction on firm body language and voice, a guide to the five principles of saying no [38] and the Alcohol Abstinence Self-Efficacy Scale [39]; the “feeling better” module includes relaxation exercises, lists of positive thoughts [40] and urge surfing [38]. In this study, we will test the app version of the intervention for the first time, targeting university students who consume alcohol over the national recommendations of no more than 9 and 14 standard drinks/week for woman and men, respectively.

**Figure 3.** Screen dumps from each of the three apps described in the article. From left to right: Promillekoll, PartyPlanner, TeleCoach.



## Assessment

The baseline, 6, 12 and 18 week follow-ups, include the Daily Drinking Questionnaire (DDQ) [28], and all 10 questions in the AUDIT [41]. The DDQ was developed in the US and was adapted for use by students by Collins et al (1985). The instrument was translated and adapted for use in Sweden by the Department of Clinical Alcohol Research at Lund University, and has been used in previous studies of Swedish students at the university [42]. The AUDIT is an effective instrument for identifying problematic alcohol consumption. The instrument was developed by the World Health Organization (WHO) and measures consumption as well as signs of harmful use and dependency related to alcohol use. The Swedish version was translated and disseminated by researchers at Karolinska Institutet [43]. The assessment package also includes information about participants' mobile phone number, gender and age, and possession of a mobile phone. All follow-ups also include questions about users experience interacting with the apps. The 18-week follow-up also includes questions regarding access, during the 18 weeks since registration, to other types of treatment such as additional sources of information, medication, speaking to counselors or using other apps than the one they are randomized to; more information on these questions is available in studies on minimal Web-based intervention conducted by our group [44,45]. Invitations to participate in follow-ups are sent via e-mail, where participants are asked to log on to a secure survey to complete assessments. Two email reminders are sent out during the one-week time period allowed for response.

## Power Calculation

Primary outcomes for both studies are quantity, frequency, number of binge drinking occasions as well as mean and peak eBAC, and proportion of individuals drinking over the recommended levels of 9 and 14 drinks per week, for women and men respectively.

For Study 1, in order to answer the questions on whether the use of a mobile phone app affects alcohol consumption compared with controls, 82 individuals need to be included in each of the three groups (two intervention and one control group, 246 individuals in total) in order to detect a difference in effect size 0.10 at 5% -level with a power of 80%. This is based on an assumed correlation between DDQ, pre- and post-measurement of at least 0.5. It is likely, however, that high power can be achieved since approximately 7,000 students will be invited to participate in the study. We expect at least 1200 students to agree to participate based on the 17% recruitment level achieved in our previous study [26], but we hope more will participate since we will be targeting first- and second-term university students in this research, in contrast to all registered students regardless of study level, in the previous study. About half of the recruited students are expected to have at least hazardous use and be eligible for randomization.

For Study 2, based on the same assumption as above, 82 individuals need to be included in each group (intervention and wait-list control group), 164 individuals in total. Based on our prior study [26] we expect that approximately 30% of the study participants will have an elevated risk consumption at 6-week follow-up, and estimates based on previous experiences of similar design [46] suggest that at least 50% will want to participate in the study, suggesting that adequate power will be available.

## Ethics and Time Plan

The project was approved by the regional ethical review board on March 19, 2014 (2014/278-31/2). The study commenced in early autumn 2014. Recruitment began during the last week of September, 2014. Participants who at 6-week follow-up showed harmful use of alcohol over 9 or 14 standard drinks per week for women and men, respectively, were offered participation in the second study. All participants were followed up at 6, 12 and 18 weeks after baseline recruitment to the project. Data collection was completed in the late spring of 2015. The results

will be analyzed during the autumn of 2015, see below for the analysis plan.

## Results

Studies 1 and 2 will be analyzed separately. For both studies, descriptive statistics will be used to describe baseline characteristics. Analysis of variance (ANOVA) will be used to identify any baseline differences in age, AUDIT, quantity, frequency, number of binge drinking occasions, as well as mean and peak eBAC between the groups. Pearson's chi-square tests will be used to determine differences between the groups in the gender distribution and the proportion of participants drinking more than the weekly recommendation.

For Study 1, all participants in both studies will be included in the analysis. Latent Markov models [47] will be used to maximize differing results for Study 1 participants not offered participation in Study 2, Study 1 participants who declined participation in Study 2, and Study 2 participants.

For Study 2, a linear mixed model analysis will be used to identify changes over time in alcohol consumption outcomes: quantity, frequency, and number of binge drinking occasions, mean eBAC and peak eBAC. These analyses will be conducted per protocol—that is, including only those participants who report using the app they were assigned to, and controlling for self-reported access to other treatment sources for problematic alcohol use during the study, and having accessed the publicly available Promillekoll app prior to the study. For comparison, intention to treat analyses will be performed with all participants randomized to experimental groups and retaining baseline values for as many participants as possible.

Descriptive statistics, ANOVA and Pearson's chi-square analyses will be performed using IBM SPSS Statistics for MacOS X, Version 22 (IBM Corp). Linear mixed model analyses will be performed using Stata 13 (StataCorp). Values for averages and standard deviations are presented to three decimal places for variables where this is necessary in order to make differences visually discernible.

## Acknowledgments

This research project was funded for further development of the PartyPlanner and TeleCoach apps by project grants from the Swedish Research Council (Ref. K2012-61X-22132-01-6), for Anne H Berman. Funding for study implementation, data analysis and writing of manuscripts for submission to international peer review, English-language scientific journals, comes from the state-run Swedish Alcohol Monopoly Research Council. Further thanks are due to Charlie Lindviken for designing and programming the original PartyPlanner app as well as the TeleCoach app.

## Authors' Contributions

Authors AHB, CA and MG conceived the study design. Authors AHB, CA, and MG revised the content of the PartyPlanner and TeleCoach apps, and MF was responsible for all technical development. MG and AHB were responsible for data collection. KS and AHB conceived the analysis plan. AHB wrote the first manuscript draft. The final draft integrated comments from KS and was written by AHB and CA.

## Discussion

### Summary

Harmful alcohol use is a significant problem in university students. Even though effective face-to-face interventions are available, most students prefer the digital interventions developed in recent years [48]. Computer use has also undergone rapid changes, and today most interaction with computers is through mobile phones [49]. Literally thousands of alcohol apps have developed for mobile phones, but the evidence for their effectiveness in reducing problematic alcohol use is lacking [21,23,27]. Our research studies seek to test the effectiveness of alcohol apps for mobile phones in reducing problematic alcohol use in university students, as well what components an efficient app actually needs to include in order to be effective.

The present research project, as well as our prior research studies [26,42], is a collaboration with the student unions at the major universities in Sweden. Our research group has over 20 years experience cooperating with student unions in alcohol research, and our overall experience is that student unions are an effective partner offering both contact information to their members, credibility, and the potential for implementing positive research findings.

The two apps our research team developed for this study are based on several years of experiences in developing and testing face-to-face brief interventions, as well as computerized brief interventions using both desktop computers and interactive voice response (IVR).

### Pros and Cons

In the present project, we decided to use Web-based apps for the two applications our research team developed. This decision has both pros and cons, as Web-based apps are cheap, easy to develop and store user data that is immediately available on our own server, lowering the risk of data losses that may occur when using regular apps run on the mobile phone itself. Regarding cons, users may perceive Web-based apps as a bit slow and sometimes absent, as they are dependent on Internet access. These cons are especially important when comparing the two Web-based apps to a regular app developed by the Swedish retailing company concerning for instance user satisfaction.

## Conflicts of Interest

None declared.

## Multimedia Appendix 1

Peer review report from funder (in Swedish).

[\[PDF File \(Adobe PDF File\), 78KB-Multimedia Appendix 1\]](#)

## References

1. Grant B, Dawson DA. Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: results from the National Longitudinal Alcohol Epidemiologic Survey. *J Subst Abuse* 1997;9:103-110. [Medline: [9494942](#)]
2. O'Neill S, Parra G, Sher KJ. Clinical relevance of heavy drinking during the college years: cross-sectional and prospective perspectives. *Psychol Addict Behav* 2001 Dec;15(4):350-359. [Medline: [11767268](#)]
3. Schuckit MA. Comorbidity between substance use disorders and psychiatric conditions. *Addiction* 2006 Sep;101 Suppl 1:76-88. [doi: [10.1111/j.1360-0443.2006.01592.x](#)] [Medline: [16930163](#)]
4. Larimer ME, Cronce JM. Identification, prevention and treatment: a review of individual-focused strategies to reduce problematic alcohol consumption by college students. *J Stud Alcohol Suppl* 2002 Mar(14):148-163. [Medline: [12022721](#)]
5. Larimer ME, Cronce JM. Identification, prevention, and treatment revisited: individual-focused college drinking prevention strategies 1999-2006. *Addict Behav* 2007 Nov;32(11):2439-2468. [doi: [10.1016/j.addbeh.2007.05.006](#)] [Medline: [17604915](#)]
6. Berglund M, Andersson C, Johnson K, Öjehagen A. Appendix 9: Limitless challenge - alcohol policy in the new age. Final report of alcohol importation investigation. In: *Prevention of hazardous drinking among young adults: a systematic literature review including meta-analyses and suggestions*. Stockholm: Swedish Government Official Reports SOU 2005; 2005.
7. National Institute on Alcohol Abuse and Alcoholism (NIAAA): Brief Interventions. *Brief Interventions*. In: *Alcohol Alert*. Washington DC: Alcohol Alert; 2005:1-8.
8. Nilsen P. Brief alcohol intervention--where to from here? Challenges remain for research and practice. *Addiction* 2010 Jun;105(6):954-959. [doi: [10.1111/j.1360-0443.2009.02779.x](#)] [Medline: [20121717](#)]
9. Cronce JM, Larimer ME. Individual-focused approaches to the prevention of college student drinking. *Alcohol Res Health* 2011;34(2):210-221 [FREE Full text] [Medline: [22330220](#)]
10. Cellucci T, Krogh J, Vik P. Help seeking for alcohol problems in a college population. *J Gen Psychol* 2006 Oct;133(4):421-433. [doi: [10.3200/GENP.133.4.421-433](#)] [Medline: [17128960](#)]
11. Buscemi J, Murphy JG, Martens MP, McDevitt-Murphy ME, Dennhardt AA, Skidmore JR. Help-seeking for alcohol-related problems in college students: correlates and preferred resources. *Psychol Addict Behav* 2010 Dec;24(4):571-580. [doi: [10.1037/a0021122](#)] [Medline: [21198220](#)]
12. Elliott J, Carey K, Bolles JR. Computer-based interventions for college drinking: a qualitative review. *Addict Behav* 2008 Aug;33(8):994-1005 [FREE Full text] [doi: [10.1016/j.addbeh.2008.03.006](#)] [Medline: [18538484](#)]
13. Bewick B, Trusler K, Barkham M, Hill A, Cahill J, Mulhern B. The effectiveness of web-based interventions designed to decrease alcohol consumption--a systematic review. *Prev Med* 2008 Jul;47(1):17-26. [doi: [10.1016/j.ypmed.2008.01.005](#)] [Medline: [18302970](#)]
14. Carey KB, Scott-Sheldon LAJ, Elliott JC, Bolles JR, Carey MP. Computer-delivered interventions to reduce college student drinking: a meta-analysis. *Addiction* 2009 Nov;104(11):1807-1819 [FREE Full text] [doi: [10.1111/j.1360-0443.2009.02691.x](#)] [Medline: [19744139](#)]
15. White A, Kavanagh D, Stallman H, Klein B, Kay-Lambkin F, Proudfoot J, et al. Online alcohol interventions: a systematic review. *J Med Internet Res* 2010;12(5):e62 [FREE Full text] [doi: [10.2196/jmir.1479](#)] [Medline: [21169175](#)]
16. Riper H, Spek V, Boon B, Conijn B, Kramer J, Martin-Abello K, et al. Effectiveness of E-self-help interventions for curbing adult problem drinking: a meta-analysis. *J Med Internet Res* 2011;13(2):e42 [FREE Full text] [doi: [10.2196/jmir.1691](#)] [Medline: [21719411](#)]
17. Carey K, Scott-Sheldon LAJ, Elliott J, Garey L, Carey MP. Face-to-face versus computer-delivered alcohol interventions for college drinkers: a meta-analytic review, 1998 to 2010. *Clin Psychol Rev* 2012 Dec;32(8):690-703 [FREE Full text] [doi: [10.1016/j.cpr.2012.08.001](#)] [Medline: [23022767](#)]
18. Moreira M, Smith L, Foxcroft D. Social norms interventions to reduce alcohol misuse in university or college students. *Cochrane Database Syst Rev* 2009(3):CD006748. [doi: [10.1002/14651858.CD006748.pub2](#)] [Medline: [19588402](#)]
19. Mattila E, Lappalainen R, Pärkkä J, Salminen J, Korhonen I. Use of a mobile phone diary for observing weight management and related behaviours. *J Telemed Telecare* 2010;16(5):260-264. [doi: [10.1258/jtt.2009.091103](#)] [Medline: [20483880](#)]
20. Ly KH, Carlbring P, Andersson G. Behavioral activation-based guided self-help treatment administered through a smartphone application: study protocol for a randomized controlled trial. *Trials* 2012;13:62 [FREE Full text] [doi: [10.1186/1745-6215-13-62](#)] [Medline: [22607302](#)]



21. Weaver E, Horyniak DR, Jenkinson R, Dietze P, Lim M. "Let's get Wasted!" and other apps: characteristics, acceptability, and use of alcohol-related smartphone applications. *JMIR Mhealth Uhealth* 2013;1(1):e9 [FREE Full text] [doi: [10.2196/mhealth.2709](https://doi.org/10.2196/mhealth.2709)] [Medline: [25100681](https://pubmed.ncbi.nlm.nih.gov/25100681/)]
22. Savic M, Best D, Rodda S, Lubman DI. Exploring the focus and experiences of smartphone applications for addiction recovery. *J Addict Dis* 2013;32(3):310-319. [doi: [10.1080/10550887.2013.824331](https://doi.org/10.1080/10550887.2013.824331)] [Medline: [24074196](https://pubmed.ncbi.nlm.nih.gov/24074196/)]
23. Crane D, Garnett C, Brown J, West R, Michie S. Behavior change techniques in popular alcohol reduction apps: content analysis. *J Med Internet Res* 2015;17(5):e118 [FREE Full text] [doi: [10.2196/jmir.4060](https://doi.org/10.2196/jmir.4060)] [Medline: [25977135](https://pubmed.ncbi.nlm.nih.gov/25977135/)]
24. DeMartini KS, Fucito LM, O'Malley SS. Novel approaches to individual alcohol interventions for heavy drinking college students and young adults. *Curr Addict Rep* 2015 Mar;2(1):47-57. [doi: [10.1007/s40429-015-0043-1](https://doi.org/10.1007/s40429-015-0043-1)] [Medline: [26258001](https://pubmed.ncbi.nlm.nih.gov/26258001/)]
25. Witkiewitz K, Desai SA, Bowen S, Leigh BC, Kirouac M, Larimer ME. Development and evaluation of a mobile intervention for heavy drinking and smoking among college students. *Psychol Addict Behav* 2014 Sep;28(3):639-650. [doi: [10.1037/a0034747](https://doi.org/10.1037/a0034747)] [Medline: [25000269](https://pubmed.ncbi.nlm.nih.gov/25000269/)]
26. Gajecki M, Berman AH, Sinadinovic K, Rosendahl I, Andersson C. Mobile phone brief intervention applications for risky alcohol use among university students: a randomized controlled study. *Addict Sci & Clin Pract* 2014;9:1-12 [FREE Full text] [doi: [10.1186/1940-0640-9-11](https://doi.org/10.1186/1940-0640-9-11)] [Medline: [24985342](https://pubmed.ncbi.nlm.nih.gov/24985342/)]
27. Cohn AM, Hunter-Reel D, Hagman BT, Mitchell J. Promoting behavior change from alcohol use through mobile technology: the future of ecological momentary assessment. *Alcohol Clin Exp Res* 2011 Dec;35(12):2209-2215 [FREE Full text] [doi: [10.1111/j.1530-0277.2011.01571.x](https://doi.org/10.1111/j.1530-0277.2011.01571.x)] [Medline: [21689119](https://pubmed.ncbi.nlm.nih.gov/21689119/)]
28. Collins RL, Parks GA, Marlatt GA. Social determinants of alcohol consumption: the effects of social interaction and model status on the self-administration of alcohol. *J Consult Clin Psychol* 1985 Apr;53(2):189-200. [Medline: [3998247](https://pubmed.ncbi.nlm.nih.gov/3998247/)]
29. Saunders JB, Aasland OG, Amundsen A, Grant M. Alcohol consumption and related problems among primary health care patients: WHO collaborative project on early detection of persons with harmful alcohol consumption--I. *Addiction* 1993 Mar;88(3):349-362. [Medline: [8461852](https://pubmed.ncbi.nlm.nih.gov/8461852/)]
30. Bland JM, Altman DG. Regression towards the mean. *BMJ* 1994 Jun 4;308(6942):1499 [FREE Full text] [Medline: [8019287](https://pubmed.ncbi.nlm.nih.gov/8019287/)]
31. Swedish Public Health Agency (Folkhälsoinstitutet). Alcohol: Knowledge base for public health policy report (Alkohol: Kunskapsunderlag för Folkhälsopolitisk rapport). Östersund: Swedish Public Health Agency; 2010.
32. Ajzen I. The theory of planned behavior. *Organizational behavior and human decision processes* 1991;50:179-211.
33. Martens MP, Ferrier AG, Cimini MD. Do protective behavioral strategies mediate the relationship between drinking motives and alcohol use in college students? *J Stud Alcohol Drugs* 2007 Jan;68(1):106-114. [Medline: [17149524](https://pubmed.ncbi.nlm.nih.gov/17149524/)]
34. Kingsbury J. The effects of social health consequence framing on drinking intentions among college students. : Society of Behavioral Medicine (SBM); 2013 Presented at: Society of Behavioral Medicine (SBM), 34th Annual Meeting and Scientific Sessions; March 20-23; San Francisco, CA.
35. Dimeff L, Baer J, Kivlahan D, Marlatt GA. Brief Alcohol Screening and Intervention for College Students (BASICS). A Harm Reduction Approach. New York, NY: The Guilford Press; 1999.
36. Fillmore MT, Jude R. Defining "binge" drinking as five drinks per occasion or drinking to a .08% BAC: which is more sensitive to risk? *Am J Addict* 2011;20(5):468-475 [FREE Full text] [doi: [10.1111/j.1521-0391.2011.00156.x](https://doi.org/10.1111/j.1521-0391.2011.00156.x)] [Medline: [21838847](https://pubmed.ncbi.nlm.nih.gov/21838847/)]
37. Gajecki M, Andersson C, Sinadinovic K, Berman AH. Interactive Voice Response (IVR) for problematic alcohol use: a three-armed randomized controlled trial. 2013 Presented at: Society of Behavioral Medicine (SBM), 34th Annual Meeting and Scientific Sessions; March 20-23; San Francisco.
38. Nielsen A. Alkoholbehandling i Praxis (Alcohol Treatment in Practice). Copenhagen: Hans Reitzel; 2008.
39. DiClemente C, Carbonari J, Montgomery R, Hughes SO. The Alcohol Abstinence Self-Efficacy scale. *J Stud Alcohol* 1994 Mar;55(2):141-148. [Medline: [8189734](https://pubmed.ncbi.nlm.nih.gov/8189734/)]
40. Berman AH, Farzanfar R, Kristiansson M, Carlbring P, Friedman RH. Design and development of a Telephone-Linked Care (TLC) system to reduce impulsivity among violent forensic outpatients and probationers. *J Med Syst* 2012 Jun;36(3):1031-1042. [doi: [10.1007/s10916-010-9565-1](https://doi.org/10.1007/s10916-010-9565-1)] [Medline: [20721686](https://pubmed.ncbi.nlm.nih.gov/20721686/)]
41. Saunders JB, Aasland OG, Babor TF, de la Fuente J R, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction* 1993 Jun;88(6):791-804. [Medline: [8329970](https://pubmed.ncbi.nlm.nih.gov/8329970/)]
42. Andersson C. Comparison of WEB and Interactive Voice Response (IVR) methods for delivering brief alcohol interventions to hazardous-drinking university students: a randomized controlled trial. *Eur Addict Res* 2015;21(5):240-252. [doi: [10.1159/000381017](https://doi.org/10.1159/000381017)] [Medline: [25967070](https://pubmed.ncbi.nlm.nih.gov/25967070/)]
43. Bergman H, Källmén H. Alcohol use among Swedes and a psychometric evaluation of the alcohol use disorders identification test. *Alcohol Alcohol* 2002;37(3):245-251 [FREE Full text] [Medline: [12003912](https://pubmed.ncbi.nlm.nih.gov/12003912/)]
44. Berman AH, Wennberg P, Sinadinovic K. Changes in mental and physical well-being among problematic alcohol and drug users in 12-month Internet-based intervention trials. *Psychol Addict Behav* 2015 Mar;29(1):97-105. [doi: [10.1037/a0038420](https://doi.org/10.1037/a0038420)] [Medline: [25664387](https://pubmed.ncbi.nlm.nih.gov/25664387/)]

45. Sinadinovic K, Wennberg P, Johansson M, Berman AH. Targeting individuals with problematic alcohol use via Web-based cognitive-behavioral self-help modules, personalized screening feedback or assessment only: a randomized controlled trial. *Eur Addict Res* 2014;20(6):305-318 [FREE Full text] [doi: [10.1159/000362406](https://doi.org/10.1159/000362406)] [Medline: [25300885](https://pubmed.ncbi.nlm.nih.gov/25300885/)]
46. Sinadinovic K, Wennberg P, Berman AH. Short-term changes in substance use among problematic alcohol and drug users from a general population sample. *IJADR* 2014 Dec 11;3(4):277-287. [doi: [10.7895/ijadr.v3i4.186](https://doi.org/10.7895/ijadr.v3i4.186)]
47. Bartolucci F, Farcomeni A, Pennoni F. Latent Markov models: a review of a general framework for the analysis of longitudinal data with covariates. *TEST* 2014 Aug 21;23(3):433-465. [doi: [10.1007/s11749-014-0381-7](https://doi.org/10.1007/s11749-014-0381-7)]
48. Kilmer J, Cronce J, Larimer ME. College student drinking research from the 1940s to the future: where we have been and where we are going. *J Stud Alcohol Drugs Suppl* 2014;75 Suppl 17:26-35 [FREE Full text] [Medline: [24565309](https://pubmed.ncbi.nlm.nih.gov/24565309/)]
49. Statistics Sweden. Privatpersoners användning av datorer och internet (Private individuals' use of computers and Internet). Stockholm: Statistics Sweden; 2014.

## Abbreviations

**AUDIT:** Alcohol Use Disorders Identification Test

**BAC:** blood alcohol concentration

**BASICS:** Brief Alcohol Screening and Intervention for College Students

**BMI:** brief motivational intervention

**DDQ:** Daily Drinking Questionnaire

**eBAC:** estimated blood alcohol concentration

**HED:** heavy episodic drinking

**IVR:** interactive voice response

**PBS:** protective behavioral strategies

**SBI:** screening with brief intervention

**WHO:** World Health Organization

*Edited by G Eysenbach; submitted 02.07.15; peer-reviewed by M Lim; comments to author 24.07.15; accepted 15.08.15; published 22.12.15*

*Please cite as:*

*Berman AH, Gajecki M, Fredriksson M, Sinadinovic K, Andersson C*

*Mobile Phone Apps for University Students With Hazardous Alcohol Use: Study Protocol for Two Consecutive Randomized Controlled Trials*

*JMIR Res Protoc* 2015;4(4):e139

URL: <http://www.researchprotocols.org/2015/4/e139/>

doi: [10.2196/resprot.4894](https://doi.org/10.2196/resprot.4894)

PMID: [26693967](https://pubmed.ncbi.nlm.nih.gov/26693967/)

©Anne H Berman, Mikael Gajecki, Morgan Fredriksson, Kristina Sinadinovic, Claes Andersson. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 22.12.2015. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.