

Protocol

A Digital Tool to Promote Alcohol and Drug Use Screening, Brief Intervention, and Referral to Treatment Skill Translation: A Mobile App Development and Randomized Controlled Trial Protocol

Derek D Satre^{1,2*}, PhD; Khanh Ly^{3*}, BS; Maria Wamsley^{3*}, MD; Alexa Curtis^{4*}, PhD; Jason Satterfield^{3*}, PhD

¹Weill Institute for Neurosciences, Department of Psychiatry, University of California, San Francisco, CA, United States

²Kaiser Permanente Northern California, Division of Research, Oakland, CA, United States

³Division of General Internal Medicine, University of California, San Francisco, CA, United States

⁴School of Nursing and Health Professions, University of San Francisco, San Francisco, CA, United States

* all authors contributed equally

Corresponding Author:

Derek D Satre, PhD

Weill Institute for Neurosciences

Department of Psychiatry

University of California

401 Parnassus Avenue, Box 0984

San Francisco, CA, 94143

United States

Phone: 1 415 476 7382

Fax: 1 415 502 2661

Email: Derek.Satre@ucsf.edu

Abstract

Background: Translation of knowledge and skills from classroom settings to clinical practice is a major challenge in healthcare training, especially for behavioral interventions. For example, screening, brief intervention, and referral to treatment (SBIRT) is a highly-promoted approach to identifying and treating individuals at risk for alcohol or drug problems, yet effective, routine use of SBIRT has lagged.

Objective: The objective of this paper is to describe the development, pilot testing, and trial protocol of a mobile app based on the theory of planned behavior (TPB) to promote SBIRT skill translation and application.

Methods: Intended for use after classroom training occurs, the mobile app has three primary functions designed to increase behavioral intent to deliver SBIRT: (1) review skills (ie, address knowledge and beliefs about SBIRT), (2) apply skills with patients (ie, build confidence and perceived behavioral control), and (3) report performance data (ie, increase accountability and social norms and/or influence). The app includes depression and anxiety screening tools due to high comorbidity with substance use. A randomized controlled trial (RCT) is in progress among health and social service learners (N=200) recruited from 3 universities and 6 different training programs in nursing, social work, internal medicine, psychiatry, and psychology. Participants are randomized to SBIRT classroom instruction alone or classroom instruction plus app access prior to beginning their field placement rotations. TPB-based data are collected via Qualtrics or via the mobile app pre-post and SBIRT utilization, weekly for 10 weeks. Key outcomes include the frequency of and self-reported confidence in delivery of SBIRT.

Results: Beta testing with advanced practice nursing students (N=22) indicated that the app and its associated assessment tools were acceptable and useful. The system usability scale (SUS) mean was 65.8 (n=19), which indicated that the SBIRT app was acceptable but could benefit from improvement. Indeed, modifications were implemented prior to starting the trial. Enrollment of trial participants began in September 2016. Results are expected by December 2017.

Conclusions: This report describes the process of TPB-based app development and testing, and the protocol for a RCT to determine the effectiveness of the app in enhancing skill translation. If effective, this approach could improve SBIRT implementation, fidelity, and clinical outcomes.

(*JMIR Res Protoc* 2017;6(4):e55) doi: [10.2196/resprot.7070](https://doi.org/10.2196/resprot.7070)

KEYWORDS

SBIRT; mobile app; digital health; training, implementation; alcohol; drug; depression; anxiety

Introduction

The transition from learning clinical skills to sustained changes in health provider behavior is a known problem. Even when evidence-based treatments are available and implementation decisions have been made, workforce development and sustained intervention delivery present formidable challenges [1]. For example, maintaining fidelity to evidence-based treatments (eg, cognitive behavioral therapy [2]), often requires strategies to support ongoing learning such as supervision and coaching [3]. Providers must learn the new skills, practice the skills, build confidence (in themselves and the intervention), and be able to change past practice norms all within an environment that supports such changes. However, these personnel-intensive strategies can be costly and time-consuming, and have limited reach due to resource constraints. Yet with effective strategies to support skill translation [4,5], behavioral healthcare providers can effectively deliver the interventions they are trained to use. Thus, a major challenge for the implementation of evidence-based behavioral practices concerns how to deliver cost-effective support for skill translation in healthcare.

Screening, brief intervention, and referral to treatment (SBIRT) for unhealthy alcohol and drug use is an important example of a widely-trained skill that has fallen short in translation [6]. SBIRT is designed to reach individuals in health or social service settings who use substances at a range of levels, including those who may not yet meet criteria for alcohol or drug use disorders. Components include screening for hazardous drinking and drug use and related problems, delivering brief motivational interviewing-based interventions for patients at low to moderate risk, and providing referrals to addiction specialty care for those with significant problems [7]. Available evidence supports the effectiveness of screening and brief intervention in addressing hazardous drinking within primary care [8-10], although evidence for effectiveness in reducing drug use is weak and trials have been mixed [11-14]. Based on the strength of this literature, national practice guidelines for SBIRT integration into primary care and other health and social service settings have been developed [15,16].

In spite of these practice recommendations and a proliferation of SBIRT training programs, optimal skill translation to direct clinical care remains unrealized. Trainees often demonstrate classroom skill proficiency yet fail to use SBIRT in subsequent clinical placements. Commonly cited barriers to translation include provider attitudes about substance use interventions, problems with knowledge recall at the point-of-care, lack of confidence, inadequate knowledge of referral resources, as well as structural barriers in clinical settings such as limited time and competing medical demands, especially in primary care [17-19]. Studies of SBIRT skill translation and implementation have found a decrease in post-training SBIRT delivery rates over time [20,21], variability in delivery rates across health disciplines [22,23], and low fidelity to screening questions [24]. Fewer than one in six Americans report being asked about or discussing their drinking with a health professional [25], and

screening is rarely conducted in US primary care settings [26] outside the Veteran's Affairs Health System [27]. Similarly, a minority of patients in mental health settings report that providers advise them to reduce hazardous drinking or drug use [28], and a recent meta-analysis demonstrated that fidelity to motivational interviewing by clinicians is often poor [29]. These findings highlight the importance of improving skill translation in real-world health and social service settings.

Digital learning tools have been incorporated into some SBIRT training programs but have not been effectively integrated with clinical care. For example, online training modules sometimes supplement didactic presentations and demonstrations, role play with feedback, and patient encounters [30-32]. With some variability, these digital health training components have been rated as relevant and useful by trainees. Outcome studies have found that such training resulted in increased confidence in SBIRT delivery and more positive attitudes towards patients who use alcohol [33]. Yet digital tools such as online learning models have not supported skill translation over time. To our knowledge, the one SBIRT mobile app that is currently available does not incorporate background materials targeted towards trainees (eg, review of prevalence of substance use and evidence for SBIRT efficacy), nor does it include detailed support in conducting screening, delivering interventions, and treatment referral resources [34]. If designed with skill translation in mind, a point-of-care mobile app with this additional content could help providers apply newly learned SBIRT skills [1].


In other health delivery contexts, mobile apps are gaining acceptance and appear to enhance training. Bullock et al (2015) found that providing a mobile app containing the Dr Companion software with 5 key medical textbooks (the iDoc app) to newly qualified doctors increased access to reference materials and was effective in supporting learning and practice [35]. The use of mobile technology, including medical apps in nursing education, has demonstrated success in improving learning outcomes and learner confidence [36]. Tablet-based patient self-administered alcohol and drug screening [37,38] and intervention [39] can increase efficiency in healthcare settings. Yet prior studies have not examined how mobile apps may enhance SBIRT training and skill translation.

Theoretical models identifying barriers and facilitators regarding learning and skill translation may help guide the development of intervention strategies to enhance skill transfer and implementation. On the level of individual provider (or learner) behavior, the theory of planned behavior (TPB) provides a well-validated, conceptual model that identifies both internal and external key factors that could influence SBIRT skill translation [40]. In the TPB model, behavioral intent (to perform the behavior of interest) is determined by attitudes and/or beliefs about the behavior, perceived social norms, and perceived behavioral control. The TPB is contextualized for SBIRT skill translation in Figure 1, allowing us to assess learners on each of these variables and to provide matched interventions as needed to promote SBIRT usage. For example, we provide

information on the extent of substance use and related problems, and SBIRT efficacy to shape attitudes regarding the value of screening and treatment. We provide information on standards of care (eg, that SBIRT has been recommended by the US Preventive Task Force and many health professional bodies) to influence perceived social norms. Moreover, we provide tools to support practicing SBIRT to positively impact both attitudes and perceived behavioral control such as trainees' confidence in successfully performing SBIRT and integrating it into clinical care.

The aim of this paper is to describe the process of TPB-based mobile alcohol and drug SBIRT app development, beta testing, and protocol for a randomized controlled trial (RCT) comparing health professional learners with access to the app (intervention arm) to learners without access to the app (control arm). We hypothesize that participants in the intervention arm will be more likely to deliver SBIRT in clinical placements than those in the control arm and will be more likely to report intention to deliver SBIRT in the future. We also hypothesize that intervention participants will report more positive beliefs about SBIRT, greater knowledge, and greater perceived control over SBIRT delivery in clinic.

Figure 1. Theory of planned behavior as applied to screening, brief intervention, and referral to treatment (SBIRT) skill translation. Model adapted from Ajzen (1991). SUD: substance use disorder; SUS: system usability scale.



Methods

App Development

An interprofessional team worked to develop the TPB-based app. The team was comprised of faculty members with prior research expertise in SBIRT training and implementation and included two clinical psychologists with doctorate degrees, an internal medicine physician, an advanced practice nurse with a

doctorate degree, and an experienced project manager. Existing mobile apps were identified to determine needs in the field and to learn about components often emphasized in enhancing behavior change, for example, mobile apps for SBIRT [34], Change Talk for Childhood Obesity [41], and Epocrates [42]. Digital product design principles were reviewed including the creation of a product vision and end goal, character sketches of potential users (“personas”), features of currently available

products, and emerging mobile app tools that simplify the user interface and promote app usage [43]. The app's purpose and vision evolved based on faculty and learner responses to initial designs. As follow-up to a needs assessment questionnaire, faculty members from different training programs were asked about the potential utility of a mobile app to increase learners' use of SBIRT. Learners identified specific features and content they wanted while using the app during clinic placements. For example, learners wanted point-of-care screening tools and SBIRT frequency of use measures integrated directly into the app.

Based on this initial feedback, the app was designed as a tool for learners to use at their clinic sites that could function as both an SBIRT information resource and as a tool to assist in skills practice and implementation. Content was designed to primarily address alcohol and drug use, but screeners for depression and anxiety, which commonly co-occur with problematic substance use, were added to broaden the scope of the app and to increase its perceived value to both learners and preceptors. A key design principle was to ensure the app fit within the clinical environment and did not disrupt other training or patient care activities. Because of data security concerns and the range of service settings and medical record systems in which trainees could use SBIRT, the app was not designed to connect with local electronic health records and does not record any protected health information. Given the expense and complexity of integrating screening into healthcare records, with which the faculty had prior experience (eg, integration of alcohol screening into electronic health records in Kaiser Permanente Northern California [22]), and the fact that not all learners are placed in settings that have electronic records, we anticipated that keeping the app separate from patient record systems would maximize learner flexibility across various clinical placements and assuage concerns about loss of patient privacy.

Flow diagrams and wireframes (page schematics and screenshots) were drafted to correspond to the key components of the app. Open Health Network app developers [44] were selected as a development partner based on their prior experience in developing mobile apps for healthcare. The wireframes were given to the developers, who provided an initial alpha version. Team members tested the alpha version individually and worked with the developers to continue refining the flow and content for subsequent beta testing.

Beta Testing


Once the final beta version of the app was developed, our team chose a small cohort of nurse practitioner learners (N=22) at the University of San Francisco to beta test the app for 3 months. Our testers included learners in an advanced practice nursing training program who were enrolled in a clinic placement. Learners completed the questionnaires and processes to be used in the larger RCT. Full TPB-based surveys (see "Pre/Post Assessment Questions") were repeated at the end of the beta-testing period, followed by a debrief focus group. The team tested and refined the app prior to starting the controlled trial. Beta testing results are described below.

Randomized Controlled Trial

Study Design

The study is a RCT of an SBIRT mobile app to facilitate skill translation from classroom to clinical placements among 200 graduate and post-graduate learners (Figure 2). Following SBIRT instruction, participants are enrolled and randomized to the experimental (use of the app) or control (no access to the app) condition. All participants complete self-report measures over the study duration (10 weeks). The trial runs from fall 2016 through spring 2017.

Figure 2. Overall study design for the development and testing of an alcohol and drug screening, brief intervention, and referral to treatment (SBIRT) mobile app to enhance skill translation. Figure adapted from the CONSORT Group.



Setting

The trial is being carried out at 3 sites, among 6 training programs: San Francisco State University (Social Work and Nursing), University of California, San Francisco (Internal Medicine and Psychiatry), and University of San Francisco (Nursing and Psychology). Institutional review boards at the 3 sites approved all study procedures. The study is not registered with ClinicalTrials.gov because the design does not include a collection of clinical data or patient-level outcomes.

Participants

Study participants (trainees, N=200) are adult health professional learners in one of the designated training programs. Participants must have had prior classroom or online training in SBIRT within the past year and may not have previously used an SBIRT app. For learners who have not yet completed classroom-based SBIRT instruction, they are required to complete the following 3 online training modules developed by the research team: (1) Introduction to SBIRT, (2) Screening, and (3) Brief Intervention. The learner must be enrolled in a field placement and is required to have a mobile device to be in the study (Android or iOS). Field placements include a range of private and public healthcare and social service agencies in the San Francisco Bay Area.

Recruitment and Randomization

Project faculty identified SBIRT educators at the participating training programs and received permission to recruit students. Students are recruited during scheduled classroom time using a detailed information sheet that specifies expectations, timing, and types of data to be collected. Absent learners are invited to

participate via email recruitment. Participants provide informed consent to participate either live in classroom settings or via email.

Standard randomization procedures used in behavioral intervention studies are followed [45] using a variable block size with a 1:1 allocation to the intervention or control arm design. These procedures are carried out by the study project manager. Learners are assigned an identification number that is used for the randomization. A Web-based randomization tool is used to generate group assignments. Randomization is stratified by training program in order to have an even distribution of learners in the intervention and control groups from each program.

Intervention Arm

Participants in the intervention group are asked to download the app, use it in their clinical rotations (either on their personal mobile phones or tablets), and complete periodic questionnaires via a Qualtrics link. The learners have the opportunity to use the app as much as they need to review SBIRT, receive guidance on structured steps in SBIRT delivery, and receive tailored recommendations on what they can do to improve. We included modest incentives for app use to maximize our ability to measure the potential effects of app use in the trial.

Control Arm

Participants in the control group do not download or use the mobile app in their practicum placements but have access to usual teaching materials and supervisors. Control participants complete periodic questionnaires via a Qualtrics link throughout

the study period. Upon completion of the study, they are invited to download and use the app at their discretion.

Data Collection

At baseline, all participants answer a TPB-based questionnaire via Qualtrics (control) or directly on the app (intervention). At the end of each week, all participants are asked to respond to a brief Qualtrics survey about how often they used SBIRT either by text message (short message service, SMS) or by email. Upon completion of their clinical rotation, all learners are asked to repeat the original TPB-based questionnaire and to provide general feedback about either the app usage (intervention) or their general satisfaction with SBIRT (control).

Incentives

Learners in both groups receive incentives for participating. The incentives are intended to enhance motivation of the learners to use the mobile app and complete study questionnaires. The learners receive Amazon gift cards throughout the study valued at US \$20 at baseline, \$2.50 for each completed SBIRT usage questionnaire, and \$20 at the end of the study for answering the final questionnaire. Maximum payment is US \$65 plus participation in a US \$50 gift card lottery based on the completion of the SBIRT usage questionnaires.

Measures

Data on participants include demographic characteristics, training institution and level of training, type of patients served in clinical placement, and pre-post TPB-based questions to capture the TPB constructs described above (Table 1).

Table 1. Measures used in the screening, brief intervention, and referral to treatment (SBIRT) mobile app randomized controlled trial (RCT).

Source	Data elements	Instrument	Timeline
All participants	Demographic characteristics	Self-report	Baseline
All participants	Type of clinical placement and population served	Self-report	Baseline
All participants	SBIRT ^a attitudes, norms, behavioral control	TPB ^b -based survey (22 items)	Baseline and 10 weeks
All participants	Delivery of SBIRT components during clinical placement	Survey	Weekly
All participants	Satisfaction with the app and usability	SUS ^c (10 items)	10 weeks
Control participants	Satisfaction with SBIRT	Survey (10 items)	10 weeks

^aSBIRT: screening, brief intervention, and referral to treatment.

^bTPB: theory of planned behavior.

^cSUS: system usability scale.

Pre-Post Assessment Questions

The team developed a 22-item questionnaire based on the TPB model. Likert-scaled items assess attitudes and beliefs including importance and efficacy of SBIRT, perceived patient willingness to participate in SBIRT, substance use epidemiology and clinical significance, and subjective norms and perceived behavioral control in the clinic setting. Three items assess confidence in the respondents' ability to screen for alcohol or drug use problems, deliver a brief intervention, and to make referrals. One item assesses intent to perform SBIRT "whenever possible in my clinical/field placement." All participants complete this questionnaire at baseline and again at 10 weeks. For intervention participants, baseline TPB responses are used to tailor their app experience by making specific recommendations of what the learner might need to review within the app's library.

The system usability scale (SUS) [46] is a 10-item Likert scale instrument developed to measure aspects of usability including system complexity and need for support and training. It yields a single score ranging from 0 to 100. Intervention group participants complete this measure at follow-up.

Satisfaction

We developed a 10-item Likert scale questionnaire to measure the experiences of control group participants, as a counterpart to the SUS. Items include barriers and challenges to SBIRT

delivery to determine why some participants might not have used SBIRT in the context of their clinical placement. We included the satisfaction questionnaire for control participants only because we want to ensure they have an equivalent number of questions to the intervention participants, and we were concerned about survey burden with our intervention participants.

Utilization.

At the end of each clinic week, every participant is sent (via email or text) a Qualtrics link asking them to report the total number of patients they have seen in the preceding week. Participants then are asked how many of those patients they screened for alcohol, drug, or tobacco use, how many they did a brief intervention with, and how many they either referred to a specialty substance use treatment clinic or discussed with their field supervisor.

Intervention


Design and Function

The SBIRT app has the following three primary functions to address TPB concepts: (1) Review SBIRT skills (to help change beliefs and attitudes), (2) Apply SBIRT skills in clinic practice (to help impact attitudes, perceived behavioral control), and (3) Report SBIRT use (to report norms, as well as study outcomes). A fourth component, the Tools section, includes additional

reference material and links (Figure 3). After downloading the app, intervention learners create an account and complete the pre-TPB survey. Immediate TPB results are given to the learners along with tailored recommendations on what they should do next. For example, if learners score low on SBIRT knowledge,

they are directed to the Review section. A progress checklist in the Tools section reminds them of “homework” they still need to complete. Throughout the study intervention learners are reminded to use the app via the weekly SBIRT usage surveys and periodic text messages promoting app usage.

Figure 3. Screen shot of the screening, brief intervention, and referral to treatment (SBIRT) mobile app home screen.



Review

The Review section includes content taught in the classroom (which is also available to learners online), as well as additional material. Subheadings include “Basics” (eg, epidemiology, drugs of abuse, consequences, defining SBIRT), “Screening” (eg, screening questions, sample scripts, and the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition [DSM-5] definition of substance use disorder [47]), “Brief Intervention”

(eg, brief advice, motivational interviewing, harm reduction), “Referral to Treatment” (eg, referral processes, pharmacotherapy), and “Key Resources” (eg, case illustrations, video demonstrations, external links to the National Institute on Alcohol Abuse and Alcoholism, and the National Institute on Drug Abuse).

Apply

The Apply section assists learners in using SBIRT while with a patient in clinic placements. This section includes screener instruments, scoring tools, scripts, and step-by-step guidance for delivering a brief intervention or referring a patient to alcohol or drug treatment. For example, the app allows users to specify what they want to screen (eg, alcohol, drugs, depression, anxiety), tailors those questions by gender and age group (18 to 64 versus 65 and up), includes single-question hazardous drinking and drug use screeners, as well as the Alcohol Use Disorders Identification Test (AUDIT) [48], CRAFFT (CRAFFT is a mnemonic acronym of first letters of key words in six screening questions) [49], and Drug Abuse Screening Test (DAST) [50]. Depression screening using the Patient Health Questionnaire (PHQ-2 and PHQ-9) [51] and anxiety screening using the Generalized Anxiety Disorder (GAD-2 and GAD-7) [52] measures are also included due to high comorbidity with substance use and commonalities in intervention approaches. Other subsections include tips for delivering brief interventions, including brief negotiated interviews/motivational interviewing, and suggestions for making referrals (eg, referral processes, lists of local treatment resources, and national treatment locators).

Report


The Report section was originally conceptualized as a library of tools for instructors and clinical preceptors to track and

evaluate their learners. We initially included a collection of pre-post surveys and weekly SBIRT usage items that intervention participants would complete. In order to standardize the data collection procedures across control and intervention participants, all weekly SBIRT usage surveys were completed via Qualtrics. Intervention participants still completed the pre-post TPB surveys and final satisfaction surveys on the app. However, instructors do not receive reports regarding use of the app or SBIRT by learners.

Tools

The Tools section includes social networking, feedback and tracking, and gamification or incentive-building tools. Social networking tools include the “social connection” to send message questions to the study team or other app users. The “Progress Checklist” allows learners to check their progress on which pages they have visited and which pages they still need to review. “Technological Support” is included in this section for those who have technical difficulties and can contact the app developers directly for help. “Leaderboard” is a page on which other learners who are using the app are listed, along with a point system indicating frequency of app use. Leaderboard rankings were tied to lottery tickets and bonus incentives (Figure 4).

Figure 4. Screenshots of the Review, Apply, Report, and Tools subsections of the screening, brief intervention, and referral to treatment (SBIRT) mobile app.



Analyses

Analyses will be conducted using SAS (SAS Institute Inc., 2011). Descriptive statistics, including distributions, means, standard deviations, skewness, and kurtosis will be obtained for all variables. Continuous measures will be tested for normality and homogeneity of variance. If the distribution is normal, Likert-scale responses will be analyzed as continuous scores [53]. Chi-square tests, student *t* tests, and analysis of variance (ANOVA) tests will be used to determine inclusion in multivariate regression models. Bivariate analyses will examine rate of SBIRT delivery in the two arms and comparison of TPB-based measures (eg, beliefs about SBIRT, social norms and influence, and perceived behavioral control). Multivariate

analyses (logistic and multiple regression) will examine the impact of these factors on SBIRT delivery.

Results

Beta Testing

Initial results focus on beta testing with student learners from the University of San Francisco School of Nursing. Beta testing was completed in summer 2016. The SUS mean was 65.8 (n=19) which indicates that the SBIRT app is acceptable but needs improvements before rolling out to a larger study sample. Debrief participants reported satisfaction with the Apply and Review sections, which included brief intervention scripts, video

demonstrations, the level of detail included in the “Referral to Treatment” section, and inclusion of the PHQ-9 (because this is often required in clinic settings to screen for depression). Suggestions for improvement focused on ease of sign-on and reducing the need for navigation (eg, by having multiple scale items appear on a single screen). These formative beta test data were used for app improvement in preparation for the RCT.

Randomized Controlled Trial

Enrollment of trial participants began in September 2016 and recruitment is ongoing. Trial results are anticipated to be available in late 2017.

Discussion

Principal Findings

The study team found that the TBP model was a useful framework for SBIRT mobile app development and that beta testers responded positively overall to the content and features of the app. The app was developed as a tool to promote translation of substance use screening and intervention skills from classroom to clinical settings. Our intent was to assist in workforce development and promote the broader use of evidence-based interventions to reduce alcohol and drug problems among patients in healthcare and social service settings. We used an app to support SBIRT skill translation, embedded in a TPB-based approach to learning, in order to inform the field regarding how mobile app technology may be used to reinforce pedagogy, improve implementation, and enhance patient care. The app, “UCSF OHN SBIRT App,” has been positively reviewed online [54] and is now publicly

available for free downloading (iOS only) via the iTunes store [55] (Figure 5). The RCT in process will determine whether the app has a significant impact on SBIRT skill translation, including rates of SBIRT delivery, learner attitudes, and intent to deliver SBIRT.

Based on the evidence and the need for intervention tools usable across settings to reduce alcohol- and drug-related problems, SBIRT instruction in both graduate training programs and continuing education settings for healthcare professionals has been spearheaded by the US Substance Abuse and Mental Health Services Administration, and training opportunities have expanded rapidly over the past 10 years. If efficacy is demonstrated, the mobile app developed by the study team may serve as a useful tool to improve training for healthcare providers and enhance patient care.

This theory-based mobile app serves as a reference guide, a clinical tool, and a data collection instrument. Learners are expected to complete the initial TPB assessment questions before starting their clinical rotations and are then asked to use the app as often as possible during the course of providing direct care. The reporting function frequency of completion is dependent on the structure of the clinical rotation and the needs of the training program and/or preceptors. Although learners’ use of SBIRT is ultimately limited by what their clinical rotation and preceptor allows, this tool may increase the likelihood of effective SBIRT delivery in healthcare and social service settings. This initial presentation describes our mobile app development process, beta testing, and randomized trial methods, which aim to determine the potential impact of this digital tool.

Figure 5. Icon of the screening, brief intervention, and referral to treatment (SBIRT) mobile app.



Limitations

The RCT is conducted in the context of graduate training in the schools in nursing, psychology, social work, and medicine, and may not generalize to other types of professional training or to providers learning SBIRT in the context of continuing education. A limitation of the trial methodology that could impact our study results is the inclusion of incentives within the intervention arm for participants to use the app, which likely would not exist in actual clinical settings. In addition, some clinical settings and supervisors may not be support the use of SBIRT, and this could impact a learner's ability to use the app. Although use of mobile devices is becoming widespread, limitations in access to technology could impact the reach of this tool [36,56]. The app was not designed to integrate responses to screening measures to electronic health records, which could limit its applicability in some clinical settings. Similarly, issues such as adherence to

app usage, appropriate use of technology in the workplace, etiquette, and distraction need to be addressed in future studies [57] to effectively integrate mobile apps into health and social service settings.

Conclusions

In behavioral health, mobile apps have primarily been directed toward patients, including alcohol and drug use reduction [58-61], smoking cessation [62], management of depression [63], and other mental health conditions [64,65]. Our approach is innovative in that it uses a skill translation theory-based intervention to target care providers and improve service delivery for important behavioral health problems. If effective, the mobile app could be scaled-up to reach a wider clinical audience and may be useful in future work on developing models of SBIRT fidelity and broader approaches to improving skill translation.

Acknowledgments

This study was supported by a grant from the Substance Abuse and Mental Health Services Administration (U79TI025404). We thank Agatha Hinman for assistance in preparing the manuscript.

Conflicts of Interest

None declared.

Multimedia Appendix 5

CONSORT eHealth checklist V1.6.1.

[\[PDF File \(Adobe PDF File\), 7MB-Multimedia Appendix 5\]](#)

References

1. Abidi L, Oenema A, Nilsen P, Anderson P, van de Mheen D. Strategies to overcome barriers to implementation of alcohol screening and brief intervention in general practice: a Delphi study among healthcare professionals and addiction prevention experts. *Prev Sci* 2016 Dec;17(6):689-699 [FREE Full text] [doi: [10.1007/s11121-016-0653-4](https://doi.org/10.1007/s11121-016-0653-4)] [Medline: [27167074](https://pubmed.ncbi.nlm.nih.gov/27167074/)]
2. Teague GB, Mueser KT, Rapp CA. Advances in fidelity measurement for mental health services research: four measures. *Psychiatr Serv* 2012 Aug;63(8):765-771 [FREE Full text] [doi: [10.1176/appi.ps.201100430](https://doi.org/10.1176/appi.ps.201100430)] [Medline: [22854723](https://pubmed.ncbi.nlm.nih.gov/22854723/)]
3. Webster-Stratton CH, Reid MJ, Marsenich L. Improving therapist fidelity during implementation of evidence-based practices: Incredible Years program. *Psychiatr Serv* 2014 Jun 01;65(6):789-795. [doi: [10.1176/appi.ps.201200177](https://doi.org/10.1176/appi.ps.201200177)] [Medline: [24686513](https://pubmed.ncbi.nlm.nih.gov/24686513/)]
4. Goldner EM, Jeffries V, Bilsker D, Jenkins E, Menear M, Petermann L. Knowledge translation in mental health: a scoping review. *Healthc Policy* 2011 Nov;7(2):83-98 [FREE Full text] [Medline: [23115572](https://pubmed.ncbi.nlm.nih.gov/23115572/)]
5. Ravitz P, Wondimagegn D, Pain C, Araya M, Alem A, Baheretibeb Y, et al. Psychotherapy knowledge translation and interpersonal psychotherapy: using best-education practices to transform mental health care in Canada and Ethiopia. *Am J Psychother* 2014;68(4):463-488. [Medline: [26453347](https://pubmed.ncbi.nlm.nih.gov/26453347/)]
6. Reho K, Agle J, DeSalle M, Gassman RA. Are we there yet? A review of screening, brief intervention, and referral to treatment (SBIRT) implementation fidelity tools and proficiency checklists. *J Prim Prev* 2016 Aug;37(4):377-388. [doi: [10.1007/s10935-016-0431-x](https://doi.org/10.1007/s10935-016-0431-x)] [Medline: [27062499](https://pubmed.ncbi.nlm.nih.gov/27062499/)]
7. Babor TF, McRee BG, Kassebaum PA, Grimaldi PL, Ahmed K, Bray J. Screening, brief intervention, and referral to treatment (SBIRT): toward a public health approach to the management of substance abuse. *Subst Abuse* 2007;28(3):7-30. [doi: [10.1300/J465v28n03_03](https://doi.org/10.1300/J465v28n03_03)] [Medline: [18077300](https://pubmed.ncbi.nlm.nih.gov/18077300/)]
8. Kaner E, Bland M, Cassidy P, Coulton S, Deluca P, Drummond C, et al. Screening and brief interventions for hazardous and harmful alcohol use in primary care: a cluster randomised controlled trial protocol. *BMC Public Health* 2009 Aug 10;9:287 [FREE Full text] [doi: [10.1186/1471-2458-9-287](https://doi.org/10.1186/1471-2458-9-287)] [Medline: [19664255](https://pubmed.ncbi.nlm.nih.gov/19664255/)]
9. Bertholet N, Daepfen J, Wietlisbach V, Fleming M, Burnand B. Reduction of alcohol consumption by brief alcohol intervention in primary care: systematic review and meta-analysis. *Arch Intern Med* 2005 May 09;165(9):986-995. [doi: [10.1001/archinte.165.9.986](https://doi.org/10.1001/archinte.165.9.986)] [Medline: [15883236](https://pubmed.ncbi.nlm.nih.gov/15883236/)]
10. O'Donnell A, Anderson P, Newbury-Birch D, Schulte B, Schmidt C, Reimer J, et al. The impact of brief alcohol interventions in primary healthcare: a systematic review of reviews. *Alcohol Alcohol* 2014;49(1):66-78 [FREE Full text] [doi: [10.1093/alcac/agt170](https://doi.org/10.1093/alcac/agt170)] [Medline: [24232177](https://pubmed.ncbi.nlm.nih.gov/24232177/)]

11. Roy-Byrne P, Bumgardner K, Krupski A, Dunn C, Ries R, Donovan D, et al. Brief intervention for problem drug use in safety-net primary care settings: a randomized clinical trial. *JAMA* 2014 Aug 06;312(5):492-501 [FREE Full text] [doi: [10.1001/jama.2014.7860](https://doi.org/10.1001/jama.2014.7860)] [Medline: [25096689](https://pubmed.ncbi.nlm.nih.gov/25096689/)]
12. Saitz R, Palfai TP, Cheng DM, Alford DP, Bernstein JA, Lloyd-Travaglini CA, et al. Screening and brief intervention for drug use in primary care: the ASPIRE randomized clinical trial. *JAMA* 2014 Aug 06;312(5):502-513 [FREE Full text] [doi: [10.1001/jama.2014.7862](https://doi.org/10.1001/jama.2014.7862)] [Medline: [25096690](https://pubmed.ncbi.nlm.nih.gov/25096690/)]
13. Aldridge A, Dowd W, Bray J. The relative impact of brief treatment versus brief intervention in primary health-care screening programs for substance use disorders. *Addiction* 2017 Feb;112 Suppl 2:54-64. [doi: [10.1111/add.13653](https://doi.org/10.1111/add.13653)] [Medline: [28074568](https://pubmed.ncbi.nlm.nih.gov/28074568/)]
14. Schwartz RP, Gryczynski J, Mitchell SG, Gonzales A, Moseley A, Peterson TR, et al. Computerized versus in-person brief intervention for drug misuse: a randomized clinical trial. *Addiction* 2014 Jul;109(7):1091-1098 [FREE Full text] [doi: [10.1111/add.12502](https://doi.org/10.1111/add.12502)] [Medline: [24520906](https://pubmed.ncbi.nlm.nih.gov/24520906/)]
15. Moyer VA, Preventive Services Task Force. Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: U.S. preventive services task force recommendation statement. *Ann Intern Med* 2013 Aug 06;159(3):210-218. [doi: [10.7326/0003-4819-159-3-201308060-00652](https://doi.org/10.7326/0003-4819-159-3-201308060-00652)] [Medline: [23698791](https://pubmed.ncbi.nlm.nih.gov/23698791/)]
16. Substance Abuse and Mental Health Services Administration. Technical Assistance Publication Series TAP 33. Rockville, MD: U.S. Department of Health and Human Services; 2013. Systems-level implementation of screening, brief intervention, and referral to treatment URL:<http://www.integration.samhsa.gov/sbirt/TAP33.pdf> [WebCite Cache ID 6pSHP7kM9]
17. Rahm AK, Boggs JM, Martin C, Price DW, Beck A, Backer TE, et al. Facilitators and barriers to implementing screening, brief intervention, and referral to treatment (SBIRT) in primary care in integrated health care settings. *Subst Abus* 2015;36(3):281-288. [doi: [10.1080/08897077.2014.951140](https://doi.org/10.1080/08897077.2014.951140)] [Medline: [25127073](https://pubmed.ncbi.nlm.nih.gov/25127073/)]
18. Satre DD, McCance-Katz EF, Moreno-John G, Julian KA, O'Sullivan PS, Satterfield JM. Using needs assessment to develop curricula for screening, brief intervention, and referral to treatment (SBIRT) in academic and community health settings. *Subst Abus* 2012;33(3):298-302 [FREE Full text] [doi: [10.1080/08897077.2011.640100](https://doi.org/10.1080/08897077.2011.640100)] [Medline: [22738009](https://pubmed.ncbi.nlm.nih.gov/22738009/)]
19. Wamsley MA, Steiger S, Julian KA, Gleason N, O'Sullivan PS, Guy M, et al. Teaching residents screening, brief intervention, and referral to treatment (SBIRT) skills for alcohol use: using chart-stimulated recall to assess curricular impact. *Subst Abus* 2016;37(3):419-426. [doi: [10.1080/08897077.2015.1127869](https://doi.org/10.1080/08897077.2015.1127869)] [Medline: [26675123](https://pubmed.ncbi.nlm.nih.gov/26675123/)]
20. van Beurden I, Anderson P, Akkermans RP, Grol RP, Wensing M, Laurant MG. Involvement of general practitioners in managing alcohol problems: a randomized controlled trial of a tailored improvement programme. *Addiction* 2012 Sep;107(9):1601-1611. [doi: [10.1111/j.1360-0443.2012.03868.x](https://doi.org/10.1111/j.1360-0443.2012.03868.x)] [Medline: [22372573](https://pubmed.ncbi.nlm.nih.gov/22372573/)]
21. Williams EC, Johnson ML, Lapham GT, Caldeiro RM, Chew L, Fletcher GS, et al. Strategies to implement alcohol screening and brief intervention in primary care settings: a structured literature review. *Psychol Addict Behav* 2011 Jun;25(2):206-214. [doi: [10.1037/a0022102](https://doi.org/10.1037/a0022102)] [Medline: [21517141](https://pubmed.ncbi.nlm.nih.gov/21517141/)]
22. Mertens JR, Chi FW, Weisner CM, Satre DD, Ross TB, Allen S, et al. Physician versus non-physician delivery of alcohol screening, brief intervention and referral to treatment in adult primary care: the ADVISE cluster randomized controlled implementation trial. *Addict Sci Clin Pract* 2015 Nov 19;10:26 [FREE Full text] [doi: [10.1186/s13722-015-0047-0](https://doi.org/10.1186/s13722-015-0047-0)] [Medline: [26585638](https://pubmed.ncbi.nlm.nih.gov/26585638/)]
23. Sterling S, Kline-Simon AH, Satre DD, Jones A, Mertens J, Wong A, et al. Implementation of screening, brief intervention, and referral to treatment for adolescents in pediatric primary care: a cluster randomized trial. *JAMA Pediatr* 2015 Nov;169(11):e153145. [doi: [10.1001/jamapediatrics.2015.3145](https://doi.org/10.1001/jamapediatrics.2015.3145)] [Medline: [26523821](https://pubmed.ncbi.nlm.nih.gov/26523821/)]
24. Bradley KA, Lapham GT, Hawkins EJ, Achtmeyer CE, Williams EC, Thomas RM, et al. Quality concerns with routine alcohol screening in VA clinical settings. *J Gen Intern Med* 2011 Mar;26(3):299-306. [doi: [10.1007/s11606-010-1509-4](https://doi.org/10.1007/s11606-010-1509-4)] [Medline: [20859699](https://pubmed.ncbi.nlm.nih.gov/20859699/)]
25. McKnight-Eily LR, Liu Y, Brewer RD, Kanny D, Lu H, Denny CH, Centers for Disease Control and Prevention (CDC). Vital signs: communication between health professionals and their patients about alcohol use--44 states and the District of Columbia, 2011. *MMWR Morb Mortal Wkly Rep* 2014 Jan 10;63(1):16-22 [FREE Full text] [Medline: [24402468](https://pubmed.ncbi.nlm.nih.gov/24402468/)]
26. Garnick DW, Horgan CM, Merrick EL, Hoyt A. Identification and treatment of mental and substance use conditions: health plans strategies. *Med Care* 2007 Nov;45(11):1060-1067. [doi: [10.1097/MLR.0b013e31812e01bb](https://doi.org/10.1097/MLR.0b013e31812e01bb)] [Medline: [18049346](https://pubmed.ncbi.nlm.nih.gov/18049346/)]
27. Bradley KA, Williams EC, Achtmeyer CE, Volpp B, Collins BJ, Kivlahan DR. Implementation of evidence-based alcohol screening in the Veterans Health Administration. *Am J Manag Care* 2006 Oct;12(10):597-606 [FREE Full text] [Medline: [17026414](https://pubmed.ncbi.nlm.nih.gov/17026414/)]
28. Satre DD, Leibowitz AS, Mertens JR, Weisner C. Advising depression patients to reduce alcohol and drug use: factors associated with provider intervention in outpatient psychiatry. *Am J Addict* 2014;23(6):570-575 [FREE Full text] [doi: [10.1111/j.1521-0391.2014.12140.x](https://doi.org/10.1111/j.1521-0391.2014.12140.x)] [Medline: [25164533](https://pubmed.ncbi.nlm.nih.gov/25164533/)]
29. Hall K, Staiger PK, Simpson A, Best D, Lubman DI. After 30 years of dissemination, have we achieved sustained practice change in motivational interviewing? *Addiction* 2016 Jul;111(7):1144-1150. [doi: [10.1111/add.13014](https://doi.org/10.1111/add.13014)] [Medline: [26216706](https://pubmed.ncbi.nlm.nih.gov/26216706/)]
30. Clemence AJ, Balkoski VI, Schaefer BM, Lee M, Bromley N, Maisonneuve IM, et al. Multispecialty screening, brief intervention, and referral to treatment (SBIRT) training in an academic medical center: resident training experience across specialties. *Subst Abus* 2016;37(2):356-363. [doi: [10.1080/08897077.2015.1082953](https://doi.org/10.1080/08897077.2015.1082953)] [Medline: [26308425](https://pubmed.ncbi.nlm.nih.gov/26308425/)]

31. Kalu N, Cain G, McLaurin-Jones T, Scott D, Kwagyan J, Fassassi C, et al. Impact of a multicomponent screening, brief intervention, and referral to treatment (SBIRT) training curriculum on a medical residency program. *Subst Abus* 2016;37(1):242-247 [FREE Full text] [doi: [10.1080/08897077.2015.1035841](https://doi.org/10.1080/08897077.2015.1035841)] [Medline: [25961140](https://pubmed.ncbi.nlm.nih.gov/25961140/)]
32. Pringle JL, Kowalchuk A, Meyers JA, Seale JP. Equipping residents to address alcohol and drug abuse: the national SBIRT residency training project. *J Grad Med Educ* 2012 Mar;4(1):58-63 [FREE Full text] [doi: [10.4300/JGME-D-11-00019.1](https://doi.org/10.4300/JGME-D-11-00019.1)] [Medline: [23451308](https://pubmed.ncbi.nlm.nih.gov/23451308/)]
33. Puskar K, Gotham HJ, Terhorst L, Hagle H, Mitchell AM, Braxter B, et al. Effects of screening, brief intervention, and referral to treatment (SBIRT) education and training on nursing students' attitudes toward working with patients who use alcohol and drugs. *Subst Abus* 2013;34(2):122-128. [doi: [10.1080/08897077.2012.715621](https://doi.org/10.1080/08897077.2012.715621)] [Medline: [23577905](https://pubmed.ncbi.nlm.nih.gov/23577905/)]
34. Bray J. iTunes Preview. 2014. SBIRT URL:<https://itunes.apple.com/us/app/sbirt/id877624835?mt=8> [accessed 2016-11-06] [WebCite Cache ID 6lpOiZ4Fn]
35. Bullock A, Dimond R, Webb K, Lovatt J, Hardyman W, Stacey M. How a mobile app supports the learning and practice of newly qualified doctors in the UK: an intervention study. *BMC Med Educ* 2015 Apr 08;15:71 [FREE Full text] [doi: [10.1186/s12909-015-0356-8](https://doi.org/10.1186/s12909-015-0356-8)] [Medline: [25889996](https://pubmed.ncbi.nlm.nih.gov/25889996/)]
36. Raman J. Mobile technology in nursing education: where do we go from here? A review of the literature. *Nurse Educ Today* 2015 May;35(5):663-672. [doi: [10.1016/j.nedt.2015.01.018](https://doi.org/10.1016/j.nedt.2015.01.018)] [Medline: [25665926](https://pubmed.ncbi.nlm.nih.gov/25665926/)]
37. McNeely J, Strauss SM, Rotrosen J, Ramautar A, Gourevitch MN. Validation of an audio computer-assisted self-interview (ACASI) version of the alcohol, smoking and substance involvement screening test (ASSIST) in primary care patients. *Addiction* 2016 Feb;111(2):233-244 [FREE Full text] [doi: [10.1111/add.13165](https://doi.org/10.1111/add.13165)] [Medline: [26360315](https://pubmed.ncbi.nlm.nih.gov/26360315/)]
38. McNeely J, Wu L, Subramaniam G, Sharma G, Cathers LA, Svikis D, et al. Performance of the Tobacco, Alcohol, Prescription Medication, and Other Substance Use (TAPS) tool for substance use screening in primary care patients. *Ann Intern Med* 2016 Nov 15;165(10):690-699. [doi: [10.7326/M16-0317](https://doi.org/10.7326/M16-0317)] [Medline: [27595276](https://pubmed.ncbi.nlm.nih.gov/27595276/)]
39. Kalkhoran S, Appelle NA, Napoles AM, Munoz RF, Lum PJ, Alvarado N, et al. Beyond the ask and advise: implementation of a computer tablet intervention to enhance provider adherence to the 5As for smoking cessation. *J Subst Abuse Treat* 2016 Jan;60:91-100 [FREE Full text] [doi: [10.1016/j.jsat.2015.05.009](https://doi.org/10.1016/j.jsat.2015.05.009)] [Medline: [26150093](https://pubmed.ncbi.nlm.nih.gov/26150093/)]
40. Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process* 1991 Dec;50(2):179-211. [doi: [10.1016/0749-5978\(91\)90020-T](https://doi.org/10.1016/0749-5978(91)90020-T)]
41. Kognito Interactive. iTunes Preview. 2016. Change talk: childhood obesity URL:<https://itunes.apple.com/us/app/change-talk-childhood-obesity/id821851796?mt=8> [accessed 2016-11-07] [WebCite Cache ID 6lpRos7YF]
42. Epocrates. URL:<http://www.epocrates.com/> [accessed 2016-11-07] [WebCite Cache ID 6lpTdcHUd]
43. Roth WR, Vilardaga R, Wolfe N, Bricker JB, McDonnell MG. Practical considerations in the design and development of smartphone apps for behavior change. *J Contextual Behav Sci* 2014 Oct 01;3(4):269-272 [FREE Full text] [doi: [10.1016/j.jcbs.2014.08.006](https://doi.org/10.1016/j.jcbs.2014.08.006)] [Medline: [25485231](https://pubmed.ncbi.nlm.nih.gov/25485231/)]
44. Open Health Network. URL:<http://www.openhealth.cc/> [accessed 2016-11-07] [WebCite Cache ID 6lpPiQUz3]
45. Hulley S, Newman T, Cummings S, Hearst N, Grady D, Browner W. *Designing Clinical Research: An Epidemiologic Approach*. 2nd ed. PA: Lippincott Williams & Wilkins; 2001.
46. usability.gov. System Usability Scale (SUS). URL:<https://www.usability.gov/how-to-and-tools/methods/system-usability-scale.html> [accessed 2016-11-06] [WebCite Cache ID 6lpNfBPGy]
47. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Arlington, VA: American Psychiatric Publishing; 2013.
48. Babor T, Higgins-Biddle J, Saunders J, Monteiro M. *AUDIT: The Alcohol Use Disorders Identification Test. Guidelines for Use in Primary Care. Second Edition*. Geneva: World Health Organization; 2001. URL:http://apps.who.int/iris/bitstream/10665/67205/1/WHO_MSD_MSB_01.6a.pdf [WebCite Cache ID 6pSKGobKg]
49. Knight JR, Sherritt L, Shrier LA, Harris SK, Chang G. Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. *Arch Pediatr Adolesc Med* 2002 Jun;156(6):607-614. [Medline: [12038895](https://pubmed.ncbi.nlm.nih.gov/12038895/)]
50. Winters KC. Development of an adolescent alcohol and other drug abuse screening scale: personal experience screening questionnaire. *Addict Behav* 1992;17(5):479-490. [Medline: [1332434](https://pubmed.ncbi.nlm.nih.gov/1332434/)]
51. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001 Sep;16(9):606-613 [FREE Full text] [Medline: [11556941](https://pubmed.ncbi.nlm.nih.gov/11556941/)]
52. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006 May 22;166(10):1092-1097. [doi: [10.1001/archinte.166.10.1092](https://doi.org/10.1001/archinte.166.10.1092)] [Medline: [16717171](https://pubmed.ncbi.nlm.nih.gov/16717171/)]
53. Sullivan GM, Artino AR. Analyzing and interpreting data from likert-type scales. *J Grad Med Educ* 2013 Dec;5(4):541-542 [FREE Full text] [doi: [10.4300/JGME-5-4-18](https://doi.org/10.4300/JGME-5-4-18)] [Medline: [24454995](https://pubmed.ncbi.nlm.nih.gov/24454995/)]
54. Maurer D. iMedicalApps. 2016. Review of UCSF OHN SBIRT app, evidence based intervention for substance abuse URL:<http://www.imedicalapps.com/2016/08/ucsf-ohn-sbirt-app-evidence-based-intervention-substance-abuse/> [accessed 2016-11-07] [WebCite Cache ID 6lpQxrAQd]
55. Open Cancer Network. iTunes Preview. 2016. OHN SBIRT URL:<https://itunes.apple.com/us/app/ohn-sbirt/id1109510478?mt=8> [accessed 2017-02-17] [WebCite Cache ID 6oLdkPyAy]

56. Muench F. The promises and pitfalls of digital technology in its application to alcohol treatment. *Alcohol Res* 2014;36(1):131-142 [[FREE Full text](#)] [Medline: [26259008](#)]
57. Bullock A, Webb K. Technology in postgraduate medical education: a dynamic influence on learning? *Postgrad Med J* 2015 Nov;91(1081):646-650 [[FREE Full text](#)] [doi: [10.1136/postgradmedj-2014-132809](#)] [Medline: [26341127](#)]
58. Barrio P, Ortega L, Bona X, Gual A. Development, validation, and implementation of an innovative mobile app for alcohol dependence management: protocol for the SIDEAL trial. *JMIR Res Protoc* 2016 Feb 17;5(1):e27 [[FREE Full text](#)] [doi: [10.2196/resprot.5002](#)] [Medline: [26888196](#)]
59. Berman AH, Gajecski M, Sinadinovic K, Andersson C. Mobile interventions targeting risky drinking among university students: a review. *Curr Addict Rep* 2016;3:166-174 [[FREE Full text](#)] [doi: [10.1007/s40429-016-0099-6](#)] [Medline: [27226948](#)]
60. Johnson K, Richards S, Chih M, Moon TJ, Curtis H, Gustafson DH. A pilot test of a mobile app for drug court participants. *Subst Abuse* 2016;10:1-7 [[FREE Full text](#)] [doi: [10.4137/SART.S33390](#)] [Medline: [26917964](#)]
61. Milward J, Khadjesari Z, Fincham-Campbell S, Deluca P, Watson R, Drummond C. User preferences for content, features, and style for an app to reduce harmful drinking in young adults: analysis of user feedback in app stores and focus group interviews. *JMIR Mhealth Uhealth* 2016 May 24;4(2):e47 [[FREE Full text](#)] [doi: [10.2196/mhealth.5242](#)] [Medline: [27220371](#)]
62. Finkelstein J, Cha EM. Using a mobile app to promote smoking cessation in hospitalized patients. *JMIR Mhealth Uhealth* 2016 May 06;4(2):e59 [[FREE Full text](#)] [doi: [10.2196/mhealth.5149](#)] [Medline: [27154792](#)]
63. Torous J, Staples P, Shanahan M, Lin C, Peck P, Keshavan M, et al. Utilizing a personal smartphone custom app to assess the Patient Health Questionnaire-9 (PHQ-9) depressive symptoms in patients with major depressive disorder. *JMIR Ment Health* 2015;2(1):e8 [[FREE Full text](#)] [doi: [10.2196/mental.3889](#)] [Medline: [26543914](#)]
64. Schlosser D, Campellone T, Kim D, Truong B, Vergani S, Ward C, et al. Feasibility of PRIME: a cognitive neuroscience-informed mobile app intervention to enhance motivated behavior and improve quality of life in recent onset schizophrenia. *JMIR Res Protoc* 2016 Apr 28;5(2):e77 [[FREE Full text](#)] [doi: [10.2196/resprot.5450](#)] [Medline: [27125771](#)]
65. Franklin JC, Fox KR, Franklin CR, Kleiman EM, Ribeiro JD, Jaroszewski AC, et al. A brief mobile app reduces nonsuicidal and suicidal self-injury: evidence from three randomized controlled trials. *J Consult Clin Psychol* 2016 Jun;84(6):544-557. [doi: [10.1037/ccp0000093](#)] [Medline: [27018530](#)]

Abbreviations

- PHQ:** Patient Health Questionnaire
RCT: randomized controlled trial
SBIRT: screening, brief intervention, and referral to treatment
SUS: system usability scale
TPB: theory of planned behavior

Edited by A Keepanasseril; submitted 29.11.16; peer-reviewed by J McNeely; comments to author 29.01.17; revised version received 18.02.17; accepted 18.02.17; published 18.04.17

Please cite as:

Satre DD, Ly K, Wamsley M, Curtis A, Satterfield J

A Digital Tool to Promote Alcohol and Drug Use Screening, Brief Intervention, and Referral to Treatment Skill Translation: A Mobile App Development and Randomized Controlled Trial Protocol

JMIR Res Protoc 2017;6(4):e55

URL: <http://www.researchprotocols.org/2017/4/e55/>

doi: [10.2196/resprot.7070](#)

PMID: [28420604](#)

©Derek D Satre, Khanh Ly, Maria Wamsley, Alexa Curtis, Jason Satterfield. Originally published in *JMIR Research Protocols* (<http://www.researchprotocols.org>), 18.04.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR Research Protocols*, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.