

Appendix IV. NHMRC Levels of Evidence (NHMRC, 2009)

Table 1 NHMRC Evidence Hierarchy: designations of 'levels of evidence' according to type of research question (including explanatory notes)

Level	Intervention ¹	Diagnostic accuracy ²	Prognosis	Aetiology ³	Screening Intervention
I ⁴	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies
II	A randomised controlled trial	A study of test accuracy with: an independent, blinded comparison with a valid reference standard, ⁵ among consecutive persons with a defined clinical presentation ⁶	A prospective cohort study ⁷	A prospective cohort study	A randomised controlled trial
III-1	A pseudorandomised controlled trial (i.e. alternate allocation or some other method)	A study of test accuracy with: an independent, blinded comparison with a valid reference standard, ⁵ among non-consecutive persons with a defined clinical presentation ⁶	All or none ⁸	All or none ⁹	A pseudorandomised controlled trial (i.e. alternate allocation or some other method)
III-2	A comparative study with concurrent controls: <ul style="list-style-type: none"> • Non-randomised, experimental trial⁹ • Cohort study • Case-control study • Interrupted time series with a control group 	A comparison with reference standard that does not meet the criteria required for Level II and III-1 evidence	Analysis of prognostic factors amongst persons in a single arm of a randomised controlled trial	A retrospective cohort study	A comparative study with concurrent controls: <ul style="list-style-type: none"> • Non-randomised, experimental trial • Cohort study • Case-control study
III-3	A comparative study without concurrent controls: <ul style="list-style-type: none"> • Historical control study • Two or more single arm study¹⁰ • Interrupted time series without a parallel control group 	Diagnostic case-control study ⁹	A retrospective cohort study	A case-control study	A comparative study without concurrent controls: <ul style="list-style-type: none"> • Historical control study • Two or more single arm study
IV	Case series with either post-test or pre-test/post-test outcomes	Study of diagnostic yield (no reference standard) ¹¹	Case series, or cohort study of persons at different stages of disease	A cross-sectional study or case series	Case series