

Table 1. Qualitative sampling matrix: patients, primary care providers, and practice staff members.

Patients	Primary care providers	Practice staff members
Psychiatric conditions: depression, anxiety, attention deficit hyperactivity syndrome, bipolar disease, use of controlled substances, substance abuse.	No contraceptive Rx^b: providers who do not provide any contraception that requires a prescription.	Director and managers: staff who have supervisory responsibilities and oversight of clinical and financial aspects of the practice (eg, clinical director, medical director, nurse manager).
Metabolic and endocrine conditions: diabetes, high blood pressure, high cholesterol, morbid obesity, thyroid problems.	Contraceptive Rx^b, No LARC^c: providers who provide contraceptive prescriptions (eg, pill, patch, ring, shot) but do not insert and remove long-acting reversible methods (LARC) ^c .	Work with PCPs^a: staff who assist PCPs during clinical visits and directly interact with patients (eg, nurses, medical assistants, licensed practical nurses).
Neurologic conditions: epilepsy, migraines, chronic pain, multiple sclerosis.	Contraceptive Rx and LARC^c: providers who provide contraceptive prescriptions (eg, pill, patch, ring, shot) AND insert and remove long-acting reversible methods (LARC) ^c .	Other services: other staff who may interact with women with chronic conditions (eg, social workers, pharmacists, behavioral counselors, complex care managers).
4.Other: clotting disorders, bleeding disorders, cancer, HIV/AIDs, rheumatologic disorders, gastrointestinal disorders.		

Notes:

^aPCPs: primary care providers.

^bRx: prescription.